



FUNDAÇÃO UNIVERSIDADE FEDERAL DE MATO GROSSO DO SUL
PROGRAMA DE PÓS-GRADUAÇÃO EM CIÊNCIAS VETERINÁRIAS
CURSO DE MESTRADO



**CARACTERIZAÇÃO DE *Escherichia coli* PRODUTORAS
DE BETA-LACTAMASES DE ESPECTRO ESTENDIDO
ISOLADAS DE BEZERROS**

YASMIN GARCIA MARANGONI

Campo Grande – MS
2023

YASMIN GARCIA MARANGONI

**CARACTERIZAÇÃO DE *Escherichia coli* PRODUTORAS
DE BETA-LACTAMASES DE ESPECTRO ESTENDIDO
ISOLADAS DE BEZERROS**

Characterization of Escherichia coli producers of extended spectrum beta-lactamases isolated from calves.

YASMIN GARCIA MARANGONI
Orientador: CÁSSIA REJANE BRITO LEAL

Dissertação apresentada ao Programa de Pós-Graduação em Ciências Veterinárias da Universidade Federal de Mato Grosso do Sul, como requisito parcial para a obtenção do título de Mestre em Ciências Veterinárias.

Campo Grande – MS
2023



Serviço Público Federal
Ministério da Educação

Fundação Universidade Federal de Mato Grosso do Sul



CERTIFICADO

Certificamos que a proposta intitulada "Caracterização epidemiológica, patológica e molecular de diarréias em bezerros nos biomas Cerrado e Pantanal sul-mato-grossense", registrada com o nº 1.134/2020, sob a responsabilidade de **CARLOS ALBERTO DO NASCIMENTO RAMOS** - que envolve a utilização de animais pertencentes ao filo Chordata, subfilo Vertebrata, para fins de pesquisa científica - encontra-se de acordo com os preceitos da Lei nº 11.794, de 8 de outubro de 2008, do Decreto nº 6.899, de 15 de julho de 2009, e com as normas editadas pelo Conselho Nacional de Controle de Experimentação Animal (CONCEA), e foi aprovada pela COMISSÃO DE ÉTICA NO USO DE ANIMAIS/CEUA DA UNIVERSIDADE FEDERAL DE MATO GROSSO DO SUL/UFMS, na 5ª reunião ordinária do dia 30/07/2020.

| | |
|-------------------------|---|
| FINALIDADE | <input type="checkbox"/> Ensino <input checked="" type="checkbox"/> Pesquisa Científica |
| Vigência da autorização | 01/09/2020 a 31/07/2023 |
| Espécie/Linhagem/Raça | Bos taurus/ Bos indicus/ Nelore e/ou cruzamentos |
| Nº de animais | 400 |
| Peso/Ideia | Não se aplica/ Até 90 dias |
| Sexo | Fêmeas e machos |
| Origem | Pantanal e Cerrado |

Fábio José Carvalho Faria
Coordenador da CEUA/UFMS
Campo Grande, 11 de agosto de 2020.



Documento assinado eletronicamente por **Fábio José Carvalho Faria**,
Presidente de Comissão, em 14/08/2020, às 10:36, conforme horário oficial
de Mato Grosso do Sul, com fundamento no art. 6º, § 1º, do Decreto nº
[8.519, de 8 de outubro de 2015](#).

A autenticidade deste documento pode ser conferida no site
https://sei.ufms.br/sei/controlador_externo.php?acao=documento_conferir&id_orgao_acesso_externo=0, informando o código
verificador 2092005 e o código CRC 5A2EBC8F.

Agradecimentos

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

Sinceros agradecimentos ao apoio da Universidade Federal de Mato Grosso do Sul – UFMS/MEC, Brasil, da Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), do Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ) e da Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia do Estado de Mato Grosso do Sul (FUNDECT).

À mesa de arguição e suplentes, por disponibilizarem tempo e paciência para este estudo.

Sinceros agradecimentos as equipes do Laboratório de Bacteriologia, FAMEZ/UFMS, Laboratório de Biologia Molecular, FAMEZ/UFMS e Laboratório de Fotônica, INFI/UFMS.

Aos professores, Cássia Rejane Brito Leal, Carlos A. do Nascimento Ramos e Cícero Cena, a técnica Lúcia Terezinha e aos alunos de graduação Carlos Eduardo Benites e Karine Dorneles.

Aos meus pais, Pedro e Elizabeth, e à Isabella, minha irmã.

Ao meu esposo, SeyedHadi.

Ao Aparecido.

Aos queridos amigos da FAMEZ e da cidade de Campo Grande, MS que me acompanharam de perto, mas também, aos queridos amigos do Irã, àqueles que me acompanharam mesmo há quilómetros de distância.

MARANGONI, Y.G. Caracterização de *Escherichia coli* produtoras de beta-lactamases de espectro estendido isoladas de bezerros. 2023. Mestrado – Programa de Pós-Graduação em Ciências Veterinárias. Faculdade de Medicina Veterinária e Zootecnia, Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, 2023.

RESUMO

A resistência antimicrobiana não é um fenômeno recente, mas é um problema crítico de saúde pública mundial. A produção de beta-lactamases de espectro estendido (ESBL), pelo grupo das Enterobacteriales, é o mecanismo de resistência bacteriana adquirida cada vez mais citado nas últimas décadas, presentes na interface humano-animal-ambiente. Este estudo teve como objetivo verificar em amostras fecais de bezerros, a presença de cepas de *Escherichia coli* diarreogênicas produtoras de ESBL, identificando os genes de resistência *bla*_{CTX-M-2}, *bla*_{CTX-M-8}, *bla*_{CTX-M-15}, *bla*_{SHV} e *bla*_{TEM} pelo método de Reação em Cadeia da Polimerase (PCR) e fenotipicamente pelo método de triagem de disco de aproximação. Também se objetivou a padronização de um método de identificação rápido, para isolados de *E. coli* multirresistentes, usando o método de Espectroscopia de Infravermelho com Transformada de Fourier (FT-IR). Duzentos e oito isolados de *E. coli* diarreogênicas foram submetidos à PCR, sendo a presença de pelo menos um dos genes citados verificada em 35,6% do total (n=74). A distribuição dos genes nas amostras positivas ocorreu da seguinte forma: 3,84% positivas para *bla*_{CTX-M-2}, 5,28% para *bla*_{SHV} e 29,3% para o gene *bla*_{TEM}. Os genes *bla*_{CTX-M-15} e *bla*_{CTX-M-8} não foram observados no estudo. Quanto à suscetibilidade aos antimicrobianos, as amostras apresentaram altas taxas de resistência ao grupo dos beta-lactâmicos, bem como para tetraciclina (80%), no qual 50,9% (106/208) foram fenotipicamente multirresistentes, mostrando que animais portadores de cepas bacterianas com a enzima ESBL, podem ser um reservatório constante de disseminação. Adicionalmente, 40 isolados de *E. coli* diarreogênicas sabidamente multirresistentes foram selecionados para serem mensurados no FT-IR, com espectros colhidos na faixa de 4000 a 700 cm⁻¹, 3000 a 2800 cm⁻¹ e 1800 a 800 cm⁻¹. O método de FT-IR para o aprendizado de máquina foi capaz de identificar os isolados de *E. coli* diarreogênicas multirresistentes com 70% de precisão no teste de validação, demonstrando que o FT-IR poderia ser uma ferramenta de triagem para estudos bacteriológicos e diagnóstico.

Palavras-chave: beta-lactâmicos, ESBL, FT-IR, pecuária e resistência à antibióticos.

MARANGONI, Y.G. Characterization of *Escherichia coli* producers of extended spectrum beta-lactamases isolated from calves. 2023. Master's degree – Postgraduate Program in Veterinary Science. Faculdade de Medicina Veterinária e Zootecnia, Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, 2023.

ABSTRACT

Antimicrobial resistance is not a recent phenomenon, but it is a critical problem of world public health. The production of extended spectrum beta-lactamase (ESBL) by the Enterobacteriales group is the increasingly cited bacterial resistance mechanism in recent decades, present in the human-animal-environment interface. This study aimed to verify in fecal calf samples, the presence of ESBL-producing *Escherichia coli* strains, identifying *bla*_{CTX-M-2}, *bla*_{CTX-M-8}, *bla*_{CTX-M-15}, *bla*_{SHV} and *bla*_{TEM} by the Polymerase Chain Reaction method (PCR) and phenotypically by the approach disc screening method. It also was an aim of a rapid identification method for multiresistant *E. coli* isolates, using the Fourier Transform Infrared Spectroscopy (FTIR) method. Diarrheagenic *E. coli* isolates, and eight isolates were subjected to PCR, the presence of at least one of the genes mentioned in 35.6% of the total (n = 74). A distributed genes from positive samples occurred as follows: 3.84% were positive for *bla*_{CTX-M-2}, 5.28% for *bla*_{SHV} gene and 29.3% for the *bla*_{TEM} gene. The *bla*_{CTX-M-15} and *bla*_{CTX-M-8} genes were not observed in the study. Regarding susceptibility to antimicrobials, the samples showed high resistance rates to the beta-lactam group, as well as for tetracycline (80%), in which 50.9% (106/208) were phenotypically multiresistant, showing that strains carriers with strains bacterial with ESBL enzyme can be a constant dissemination reservoir. Additionally, 40 isolates of *E. coli* diarrheagenic multiresistant were selected to be measured in the FTIR, with spectra harvested from 4000 to 700 cm⁻¹, 3000 to 2800 cm⁻¹, and 1800 to 800 cm⁻¹. The FTIR method for machine learning was able to identify the isolates of *E. coli* diarrheagenic multi-resistant with 70% accuracy in validation tests, demonstrating that the FTIR could be a screening tool for bacteriological and diagnostic studies.

Keywords: beta-lactams, ESBL, FTIR, livestock and antibiotic resistance.

LISTA DE FIGURAS

| | | |
|-------------------|---|----|
| Figura 1. | Esquema bioquímico demonstrando a ação da enzima betalactamase na ligação amina do anel betalactâmico..... | 06 |
| Figura 2. | Ilustração da posição ideal das cefalosporinas, como cefotaxima (CTX), ceftazidima (CAZ) e ceftriaxona (CRO), o monobactâmico aztronam (ATM), e o inibidor de betalactamase amoxicilina + ácido clavulônico (AMC) no teste de triagem de disco de aproximação, padronizado pelo CLSI (2008) | 07 |
| Figura 3 | Ilustração básica da funcionalidade do espectrômetro, demonstrando a (a) fonte de luz, (b) interação da luz com a amostra analisada, (c) transmissão para um fotodetector, (d) registro do espectro da absorção de radiação eletromagnética, ou seja, "impressão digital. | 10 |
| CAPÍTULO 1 | | |
| Figura 1 | Positive approximation disk screening technique demonstrating the phantom zone or absence of inhibition halo (red arrow) to verify samples possibly producing the ESBL enzyme..... | 22 |
| Figura 2 | Electrophoresis of positive samples on a 1% agarose gel, using a molecular marker of 100 base pairs, visualizing the amplification of the <i>bla</i> _{CTX-M-2} and <i>bla</i> _{SHV} (A), as well as the <i>bla</i> _{TEM} gene (B)..... | 23 |
| Figura 3 | Map of the state of Mato Grosso do Sul, Brazil, showing the positive municipalities (yellow) for the genotypic identification of the ESBL-encoding genes: <i>bla</i> _{CTX-M-2} (triangle), <i>bla</i> _{SHV} (square) and <i>bla</i> _{TEM} (circle)..... | 24 |
| CAPÍTULO 2 | | |
| Figura 1 | Experimental scheme describing the main steps from the sample (<i>E. coli</i>) preparation to FTIR spectra acquisition..... | 40 |
| Figura 2 | Average FTIR-SNV spectra for (a) <i>E. coli</i> ATCC (blue line), and (b) multidrug-resistant (MR) <i>E. coli</i> (red line) isolates. The main vibrational assignments are indicated in detail and their related molecular groups (proteins, lipids, carbohydrates, and fatty acids) are identified by colors. The symbol δ indicates the deformation vibrations. | 43 |
| Figura 3 | Principal Component Analysis for <i>E. coli</i> ATCC (blue circles/stars), and multidrug-resistant (MR) <i>E. coli</i> (red circles/stars) isolates average FTIR-SNV spectra. The score plot and respective loading for the three different ranges were analyzed: (a-b) 4000 to 800 cm ⁻¹ ; (c-d) 3000 to 2800 cm ⁻¹ ; and (e-f) 1800 to 800 cm ⁻¹ . The circles represent 70% of the sample data set used to build the predicting model, and the stars represent 30% of the sample data set used for external validation..... | 44 |

| | | |
|----------|--|----|
| Figura 4 | Violin plot for nine main PCs normalized scores selected by RFE in 1800 to 800 cm ⁻¹ range from <i>E. coli</i> ATCC® 25922 (blue), and multidrug-resistant (MR) <i>E. coli</i> (red) isolates. The thick dashed line represents the data distribution median. The data variance percentage of each PC is described above each plot..... | 46 |
| Figura 5 | (a) score plot clustering improvement due to PCs selection by RFE., and (b) loading for PC4, PC17, and PC28 from <i>E. coli</i> ATCC® 25922 (blue square), and (b) multidrug-resistant (MR) <i>E. coli</i> (red circles) isolates. | 47 |
| Figura 6 | (a) Overall accuracy obtained in the LOOCV tests for Discriminant analysis (DA), K-nearest neighbor (KNN), and Support vector machine (SVM) algorithms. Different ranges were analyzed with respective RFE-PCs: (i) 4000 to 800 cm ⁻¹ (gray bar color with right inclined line pattern) using 5 PCs; (ii) 3000 to 2800 cm ⁻¹ (light blue color with horizontal line pattern) by using 5 PCs; (iii) 1800 to 800 cm ⁻¹ (purple color with left inclined line pattern) by using 9 PCs. The highest overall accuracy was 85% for Linear SVM at (4000 to 800 cm ⁻¹) and (1800 to 800 cm ⁻¹) ranges, and Weighted KNN at (4000 to 800 cm ⁻¹) range. Confusion matrix for the performance of Linear SVM algorithm in 1800 to 800 cm ⁻¹ range by using 9 PCs (b) LOOCV test with 70% data set, and (c) external validation test with 30% data set. The input PCA data from average FTIR-SNV spectra of <i>E. coli</i> ATCC® 25922, and multidrug-resistant (MR) <i>E. coli</i> isolates..... | 47 |

LISTA DE TABELAS

| | | |
|-------------------|---|----|
| CAPÍTULO 1 | | |
| Tabela 1 | Agentes bacterianos e suas respectivas resistências antimicrobianas na medicina humana, segundo a Organização Mundial da Saúde (2017) | 04 |
| Tabela 1 | Initiators to identify the genes encoding ESBL..... | 21 |
| Tabela 2 | Correlation of the frequency of antimicrobial resistance to beta-lactams (A) by the disk diffusion method and antimicrobials from the classes of aminoglycosides, fluoroquinolones, tetracyclines, amphenicols and sulfonamides (B), with the location of the <i>bla</i> _{CTX-M- 2} genes, <i>bla</i> _{SHV} and <i>bla</i> _{TEM} from diarrheagenic <i>E. coli</i> isolates by the PCR method..... | 24 |
| Tabela 3 | Frequency of in vitro antimicrobial resistance in 208 diarrheagenic <i>Escherichia coli</i> isolates from stool and rectal swab samples..... | 25 |

SUMÁRIO

| | |
|---|----|
| RESUMO..... | i |
| ABSTRACT..... | ii |
| CAPÍTULO 1..... | 01 |
| 1 INTRODUÇÃO GERAL..... | 01 |
| 1.1. Objetivo geral..... | 02 |
| 1.2. Objetivos específicos..... | 02 |
| 2 REVISÃO DE LITERATURA..... | 03 |
| 2.1. Resistência antimicrobiana..... | 03 |
| 2.2. Espectroscopia óptica..... | 08 |
| 2.2.1. Espectroscopia no Infravermelho com Transformada de Fourier (FT-IR) | 08 |
| 3 REFERÊNCIAS BIBLIOGRÁFICAS..... | 11 |
| CAPÍTULO 2..... | 17 |
| 4 (ARTIGO I). Identificação fenotípica e genotípica de <i>Escherichia coli</i> produtoras de beta-lactamases de espectro estendido (ESBL) isoladas de fezes de bezerros..... | 17 |
| 5 (ARTIGO II). Identification of multi-resistant diarrheagenic <i>Escherichia coli</i> by using FTIR Spectroscopy and Machine Learning: a feasible strategy to improve the group classification..... | 37 |

CAPÍTULO 1

1. INTRODUÇÃO GERAL

A identificação de bactérias multirresistentes e seus genes de virulência na pecuária tem grande importância em saúde pública mundial, assim como a identificação de possíveis genes de resistência à antimicrobianos (AGRs) (HAN et al., 2017).

O rebanho comercial brasileiro cresce com taxas altas, chegando a 218,2 milhões de cabeças de gado. A região Centro-Oeste, respondeu por mais de 75 milhões dessas cabeças. Em termos municipais, Corumbá, Mato Grosso do Sul, segue em segunda posição com o maior rebanho nacional (IBGE, 2020).

A enterite em bezerros, provocada pela bactéria *Escherichia coli* tem sido relatada desde o início do século XX como um dos principais agentes envolvidos nas diarreias de origem infecciosa. Considerando que o estado de Mato Grosso do Sul é uma importante região de pecuária de corte, este agente etiológico e suas cepas diarréogênicas passaram a ser estudadas e identificadas no estado (SMITH; ORCUTT, 1925; TUTIJA et al., 2022).

A Organização Mundial da Saúde, OMS (2017) lista agentes bacterianos e suas respectivas resistências antimicrobianas na medicina humana, como um alerta à população mundial a crescente resistência global aos medicamentos antimicrobianos, entretanto, dentro da medicina veterinária já podemos observar a cosmopolitização destes AGRs listados pela OMS, como a produção de beta-lactamase de espectro estendido (ESBL) pela bactéria *E. coli* na interface humano-animal-ambiente (MELO et al., 2018; PALMEIRA et al., 2020).

Atualmente, por não existirem programas de vigilância epidemiológica de abrangência nacional ativos referentes à resistência bacteriana e a seus mecanismos, torna-se difícil estimar a proporção de cepas multirresistentes e aquelas produtoras de ESBL na bovinocultura (SANTAJIT; INDRAWATTANA, 2016; CHONG et al., 2018).

Com o surgimento de ferramentas da biologia molecular, podemos realizar análises detalhadas das sequências de ácidos nucleicos para identificação de AGRs, porém sua abrangência, seu poder discriminatório e simplicidade são variáveis e, para muitas análises, o alto custo e o tempo dificultam sua implementação na rotina laboratorial (SABAT et al., 2013).

35 Hoje é altamente desejável a busca pela implementação de novos métodos
36 para detectar microrganismos e AGRs, com resultados rápidos, reduzido custo
37 com consumíveis e fácil portabilidade (BENGTSSON-PALME et al., 2017).

38 Os estudos de espectroscopia óptica, como a Espectroscopia no
39 Infravermelho com Transformada de Fourier (FT-IR), representam uma
40 ferramenta poderosa para atingir esses objetivos, bem como em estudo futuros,
41 nos processos de vigilância epidemiológica nacional referentes à resistência
42 bacteriana e a seus mecanismos, usando abordagens baseadas em análises
43 vibracionais em conjunto com a bioinformática (NOVAIS et al., 2019).

44 Diante deste contexto relatado, este trabalho objetivou estudar a presença de
45 cepas de *E. coli* diarreogênicas produtoras de ESBL no rebanho do estado de
46 Mato Grosso do Sul, Brasil, em conjunto com o aperfeiçoamento do diagnóstico
47 de resistência antimicrobiana por técnicas de fotônica.

48

49 **1.1 OBJETIVO GERAL**

50 O presente estudo teve por objetivo investigar fenotipicamente e
51 genotipicamente a presença de linhagens de *E. coli* produtoras de ESBL das
52 famílias CTX-M (*bla*_{CTX-M-2}, *bla*_{CTX-M-8} e *bla*_{CTX-M-15}), SHV (*bla*_{SHV}) e TEM (*bla*_{TEM})
53 na microbiota intestinal de bezerros, provenientes de fazendas das quatro
54 macrorregiões do estado de Mato Grosso do Sul, Brasil, bem como realizar a
55 padronização para o método de FT-IR em isolados de *E. coli* multirresistentes.

56

57 **1.2 OBJETIVOS ESPECÍFICOS**

- 58 • Avaliar *in vitro* o perfil de suscetibilidade aos antimicrobianos em isolados
59 de *E. coli* diarreogênicas pelo método de disco difusão;
- 60 • Identificar fenotipicamente a presença de estirpes produtoras de ESBL;
- 61 • Identificar genotipicamente por meio da técnica de Reação em Cadeia da
62 Polimerase (PCR), os genes codificadores de ESBL: *bla*_{CTX-M-2}, *bla*_{CTX-M-8},
- 63 *bla*_{CTX-M-15}, *bla*_{SHV} e *bla*_{TEM};
- 64 • Avaliar o desempenho da técnica de Espetroscopia de Infravermelho por
65 Transformada de Fourier (FT-IR) para identificar cepas multirresistentes
66 de *E. coli*.

67

68 2 REVISÃO DE LITERATURA

69

70 2.1. Resistência antimicrobiana

71 A introdução de antibióticos no uso clínico foi indiscutivelmente o maior
72 avanço médico do século 20 (KATZ; BALTZ, 2016).

73 Em 1904, Paul Ehrlich e Alexander Fleming iniciaram a moderna “era dos
74 antibióticos”. Ehrlich iniciou um programa de rastreamento sistemático para
75 encontrar um medicamento contra a sífilis, baseadas em arsênico sintético
76 salvarsan, porém os primeiros relatos de resistência ao salvarsan foram em 1930
77 (EHRLICH, 1913).

78 Salvarsan foi substituído pela sulfonamida Prontosil (OTTEN, 1986),
79 considerado o primeiro antimicrobiano de amplo espectro com uso clínico,
80 entretanto foi amplamente substituído pela descoberta da penicilina por
81 Alexander Fleming em 1928 (FLEMING, 1929). A partir de 1945, a penicilina foi
82 purificada por um grupo de pesquisadores da Universidade de Oxford, e após
83 esse período, houve relatos crescentes de resistência a este medicamento
84 (ABRAHAM et al., 1941).

85 O início da “era de ouro” da descoberta de antibióticos, ocorreu entre as
86 décadas de 1940 e 1960. A maioria desses compostos ainda está em uso clínico,
87 mas sua eficácia foi prejudicada pelo aumento da resistência microbiana. Na
88 verdade, a descoberta rápida e relativamente fácil de várias classes de
89 antibióticos durante um período relativamente curto, levou ao uso excessivo
90 desses medicamentos (KATZ; BALTZ, 2016).

91 Existem diferentes mecanismos de resistência antimicrobiana: a resistência
92 intrínseca e a resistência adquirida. A resistência intrínseca ocorre de forma
93 natural, como parte de um processo de evolução bacteriana. Alguns exemplos
94 são membros da ordem Enterobacteriales, na qual a bactéria *E. coli* está
95 inserida, resistentes as classes dos macrolídeos, glicopeptídeos, bem como ao
96 ácido fusídico, linezolida e a penicilina G, considerados antimicrobianos de
97 grande uso na rotina clínica (WHO, 2012; EUCAST, 2020; CLSI, 2021).

98 Com frequência as bactérias utilizam mais de uma estratégia para evitar a
99 ação dos antimicrobianos. Na resistência adquirida, a bactéria altera seu material
100 genético de duas formas: pela mutação no seu material genético; ou introdução
101 de um material genético estranho, como genes de resistência, que podem ser

102 transferidos entre gêneros ou espécies diferentes de bactérias, principalmente
 103 devido à pressão seletiva exercida pelo uso indiscriminado de antimicrobianos
 104 (LIVERMORE et al., 2001; WHO, 2012; EUCAST, 2020).

105 O relatório O'Neill (2016) previu que, sem ação urgente, dez milhões de
 106 pessoas, por ano morrerão de infecções causadas por bactérias multirresistentes
 107 à antimicrobianos até 2050.

108 Como um alerta à população mundial a crescente resistência aos
 109 medicamentos antimicrobianos, a Organização Mundial da Saúde (OMS), lista
 110 os agentes bacterianos e suas respectivas resistências antimicrobianas (Tabela
 111 1), utilizando critérios para a seleção das bactérias, como: a) quão letal as
 112 infecções que esses microrganismos causam são; b) se o seu tratamento requer
 113 longas estadias hospitalares; d) como eles se espalham facilmente entre os
 114 animais, dos animais aos seres humanos, e de pessoa para pessoa; e) se os
 115 microrganismos podem ser prevenidos; f) quantas opções de tratamento ainda
 116 existem para esses microrganismos (WHO, 2017).

117

118 Tabela 1: Agentes bacterianos e suas respectivas resistências antimicrobianas na medicina
 119 humana, segundo a Organização Mundial da Saúde (2017).

| Prioridade crítica | Prioridade alta | Prioridade média |
|---|--|--|
| <i>Acinetobacter baumannii</i> resistente à carbapenêmicos | <i>Enterococcus faecium</i> resistente à vancomicina | <i>Streptococcus pneumoniae</i> não penicilina suscetível |
| <i>Pseudomonas aeruginosa</i> resistente à carbapenêmicos | <i>Helicobacter pylori</i> resistentes à claritromicina | <i>Haemophilus influenzae</i> resistentes à ampicilina |
| <i>Enterobacteriaceae</i> resistentes à 3ª geração de cefalosporinas e carbapenêmicos. | <i>Salmonella</i> sp resistentes à fluoroquinolonas | <i>Shigella</i> sp resistentes à fluoroquinolona |
| | <i>Staphylococcus aureus</i> resistente à vancomicina e meticilina | |
| | <i>Campylobacter</i> sp resistentes à fluoroquinolona | |
| | <i>Neisseria gonorrhoeae</i> resistentes à 3ª geração de cefalosporinas e fluoroquinolona | |

120 Fonte: WHO (2017).

121

122 Dentro da medicina veterinária já podemos observar a cosmopolização de *A.*
 123 *baumannii* (POMBA et al., 2015; EWERS et al., 2017), *P. aeruginosa*
 124 (FERNANDES et al., 2018; ELSHAFIEE et al., 2019), *Staphylococcus aureus*

125 (OLIVEIRA et al., 2016); *Salmonellas* species (COLOBATIU et al., 2015; PRIBUL
126 et al., 2017), *Campylobacter* species (FRASAO et al., 2015), bem como algumas
127 bactérias membros do grupo Enterobacteriales, como a *E. coli* (MELO et al.,
128 2018; PALMEIRA et al., 2020) na interface humano-animal-ambiente.

129 A bactéria *E. coli* é considerada comensal uma vez que, apenas uma
130 pequena parte das cepas apresentam patogenicidade responsável por
131 enfermidades. A diferenciação de cepas patogênicas (diarreogênicas) de cepas
132 não patogênicas, é baseada na produção de fatores de virulência, sendo cinco
133 os principais patótipos de *E. coli* diarreogênica para bezerros: *E. coli*
134 enterotoxigênica (ETEC), *E. coli* enteropatogênica (EPEC), *E. coli*
135 enterohemorrágica (EHEC), *E. coli* produtora de toxina Shiga (STEC), *E. coli*
136 necrotoxigênica (NTEC). Cada categoria possui capacidade de causar síndrome
137 clínica, com características epidemiológicas e patológicas específicas
138 (BLANCHARD, 2012; COURA et al., 2015).

139 Quando há disseminação de cepas de *E. coli* diarreogênicas no rebanho
140 bovino, podem surgir cepas bacterianas capazes de expressarem enzimas que
141 conferem resistência a múltiplos antimicrobianos, em resposta ao amplo uso
142 destas moléculas para tratar os casos de diarreia (BLANCHARD, 2012; COURA
143 et al., 2015; POIREL et al., 2018).

144 A classe de antimicrobianos beta-lactâmicos constitui a primeira via de
145 escolha para o tratamento de diarreia, devido a sua boa eficácia terapêutica e
146 baixa toxicidade. Representantes dessa classe são o grupo das penicilinas
147 (amoxicilina, associada ou não à clavulanato de potássio e ampicilina) e
148 cefalosporinas (1^a e 3^a geração), ambos com a presença do anel beta-lactâmico
149 em sua estrutura molecular (CONSTABLE, 2004; SPINOSA et al., 2014).


150 O mecanismo principal de ação dos beta-lactâmicos é a inibição da última
151 etapa de síntese de peptídeoglicano para a formação da parede celular
152 bacteriana, logo estes antimicrobianos somente são capazes de atuar em
153 bactérias durante a fase de crescimento logarítmica, quando, então, há
154 necessidade da síntese da parede celular (SPINOSA et al., 2014; ANDRADE;
155 DARINI, 2016).

156 O mecanismo de resistência mais importante para os beta-lactâmicos é a
157 produção de betalactamases pelas bactérias. As betalactamases produzidas por
158 diferentes bactérias possuem propriedades físicas, químicas e funcionais

variadas, sendo algumas betalactamases específicas para penicilinas (penicilinases), outras para cefalosporinas (cefaloparinases) e outras que podem atuar em ambos os grupos de antimicrobianos, como as beta-lactamases de espectro estendido (ESBL) (SPINOSA et. al, 2014).

Diversos trabalhos têm apontado para o crescimento mundial das ESBL em bactérias Gram-negativas, como membros do grupo Enterobacteriales (ELLER et al., 2014; ROCHA et al., 2016; GUNDTRAN et al., 2019; SUKMAWINATA et al., 2020; WIDODO et al., 2020). Estas enzimas quando produzidas pelas bactérias Gram-negativas são secretadas no espaço periplasmático (SPINOSA et. al, 2014) podendo hidrolisar o anel beta-lactâmico irreversivelmente da ligação amina do anel beta-lactâmico (Figura 1), impedindo a ação dos beta-lactâmicos na síntese da parede celular bacteriana (BUSH; JACOBY, 2010), hidrolisando todas as penicilinas, cefalosporinas, incluindo as de terceira e quarta geração, e o aztreonam (COQUE et al., 2008).

173



174

Figura 1: Esquema bioquímico demonstrando a ação da enzima beta-lactamase na ligação amina do anel beta-lactâmico. Arquivo pessoal.

177

A presença de ESBL tem um significado clínico importante, pois são codificadas por plasmídeo, e estes frequentemente carregam genes com co-resistência a outras classes de antimicrobianos, como, por exemplo, tetraciclinas, aminoglicosídeos e fluoroquinolonas. Portanto, as opções de antimicrobianos no tratamento de organismos produtores de ESBL são extremamente limitadas (PATERSON; BONOMO, 2005; ADEYANKINNU et al., 2014).


Dentre as variantes enzimáticas descritas, as enzimas da família SHV e TEM foram inicialmente reconhecidas em diferentes espécies bacterianas Gram-negativas, descritas em humanos a partir da década de 1960 (CANTÓN et al.,

188 2012), porém as enzimas da família CTX-M têm despertado atenção por sua alta
 189 incidência e grande capacidade de propagação (PITOOUT et al., 2005; WIDODO
 190 et al., 2020).

191 As fezes de animais de produção, que possuem cepas com ARGs, como a
 192 enzima ESBL, podem ser reservatórios constantes de disseminação aos
 193 humanos (CARATTOLI, 2008; SCHMIDT et al., 2013), ao meio ambiente e a
 194 outros animais (BOONYASIRI et al., 2014; TEKINER; OZPINAR, 2016), levando
 195 à atrasos na terapia apropriada, aumento da morbidade, mortalidade, e
 196 consequentemente, em perdas econômicas aos produtores (SANTAJIT;
 197 INDRAWATTANA, 2016; CHONG et al., 2018).

198 Padrões de resistência entre bactérias têm sido estudados utilizando o cultivo
 199 a partir de métodos fenotípicos, como o teste de triagem de disco de
 200 aproximação (JARLIER et al., 1988) (Figura 2), bem como métodos moleculares,
 201 como a Reação em Cadeia da Polimerase (PCR), localizando os genes *bla*_{CTX-}
 202 _M, *bla*_{SHV} e *bla*_{TEM} (PATERSON; BONOMO, 2005).

203



204

205 Figura 2: Ilustração da posição ideal das cefalosporinas, como cefotaxima (CTX), ceftazidima
 206 (CAZ) e ceftriaxona (CRO), o monobactâmico aztronam (ATM), e o inibidor de beta-lactamase
 207 amoxicilina + ácido clavulônico (AMC) no teste de triagem de disco de aproximação, padronizado
 208 pelo CLSI (2021). Arquivo pessoal.

209

210 Entretanto, hoje é altamente desejável a busca pela implementação de novos
 211 métodos de identificação bacteriana, dos quais não utilizem uma série de testes

212 para a identificação definitiva do organismo. Uma abordagem diferente para a
213 identificação rápida de microrganismos é baseada em técnicas espectroscópicas
214 (MAQUELIN et al., 2002).

215

216 **2.2 Espectroscopia óptica**

217 Do ponto de vista histórico, entre 1665 e 1666, Isaac Newton, um físico inglês,
218 passou a introduzir a espectroscopia ao observar a luz do sol. A luz ao passar
219 de meio de propagação para outro, como um prisma de vidro, poderia ser
220 decomposta em diversas cores (NEWTON, 1972).

221 No ano de 1802, o interesse pela espectroscopia aumentou
222 consideravelmente, quando um grupo passou a observar que substâncias
223 emitiam cores diferentes quando colocadas sob a chama, com o objetivo de
224 identificar os elementos químicos presentes. A partir de 1859, os químicos
225 Robert Wilhelm Bunsen e Gustav Kirchhoff tiveram um papel fundamental na
226 construção do primeiro espectroscópio, descobrindo que cada elemento químico
227 produz um padrão único de linhas espectrais, distribuída ao longo dos
228 comprimentos de onda (BUNSEN; KIRCHHOFF, 1862).

229 Esses e muitos estudos posteriores abriram caminhos para a utilização dessa
230 metodologia para a análise de substâncias, como a descoberta do cloreto de
231 césio e o cloreto de rubídio, bem como para a identificação de bactérias em
232 diferentes níveis taxonômicos, desde gêneros, espécies a sorotipo/sorogrupo
233 (BUNSEN; KIRCHHOFF, 1862; HELM et al. 1991; MAQUELIN et al., 2002).

234 Durante a década de 1990, houve novos desenvolvimentos de
235 equipamentos, procedimentos experimentais e análise de dados em software, o
236 que impulsionaram a novas publicações sobre a técnica de espectroscopia no
237 infravermelho com transformada de Fourier (FT-IR) para identificação bacteriana
238 (HELM et al. 1991).

239

240 **2.2.1 Espectroscopia no Infravermelho com Transformada de Fourier 241 (FT-IR).**

242 Técnicas espectroscópicas destrutivas, como a Espectrometria de Massa
243 com Ionização por Dessorção a Laser Assistida por Matriz (MALDI-TOF MS) tem
244 mostrado resultados promissores para a identificação de grupos bacterianos
245 Gram-negativos e Gram-positivos, porém inconsistências entre os protocolos


246 experimentais ou de análise de dados adotados podem dificultar o alcance de
247 reprodutibilidade, padronização e, principalmente, a portabilidade para
248 laboratórios de rotina de microbiologia clínica (SAUGET et al., 2017).

249 Considerando o bom desempenho em diferenciação de cepas, alto
250 rendimento, custo-benefício, mínimo tempo de trabalho, e fácil portabilidade,
251 métodos não destrutivos - àqueles que deixam a amostra intacta durante a
252 análise - podem ser desenvolvidos com base em espectroscopias vibracionais,
253 como o FT-IR (MAQUELIN et al., 2002; NOVAIS et al., 2019).

254 Na região do infravermelho (100 a 10000 cm⁻¹), ocorre a absorção da
255 radiação infravermelha por um determinado tipo de amostra como, por exemplo,
256 células bacterianas, causando excitação e vibração dos seus diferentes
257 compostos químicos, com o objetivo de determinar a composição molecular
258 (ácidos nucleicos, proteínas, lipídios e carboidratos) destas células bacterianas
259 como um todo, semelhantes a impressões digitais ou espectro que podem ser
260 utilizados para identificar a identidade do microrganismo (HELM et al. 1991;
261 SETTLE, 1997; STUART, 2004).

262 O método de FT-IR, utiliza um equipamento chamado de espetrômetro
263 (Figura 3). No espetrômetro, a fonte de luz emite um feixe de luz com energia e
264 comprimento de onda bem definido. Este feixe de luz ao passar pela amostra
265 analisada pode interagir com o meio e ser transmitido para um fotodetector.
266 Logo, um espectro característico da absorção de radiação eletromagnética da
267 amostra é obtido, e por meio de cálculos matemáticos, da transformada de
268 Fourier, a distância do comprimento óptico pode ser convertida para o valor da
269 frequência de radiação e vice-versa, garantindo a otimização das funções da
270 espectroscopia no infravermelho, permitindo maior sensibilidade e velocidade de
271 análise (STUART, 2004).

272



273

274 Figura 3: Ilustração básica da funcionalidade do espectrômetro, demonstrando a (a) fonte de luz,
 275 (b) interação da luz com a amostra analisada, (c) transmissão para um fotodetector, (d) registro
 276 do espectro da absorção de radiação eletromagnética, ou seja, “impressão digital. Arquivo
 277 pessoal.

278

279 O potencial do método de FT-IR para diferenciar e identificar grupos de cepas
 280 dentro de espécies Gram-positivas, como *Staphylococcus aureus* e Gram-
 281 negativas, principalmente em *E. coli*, sendo explorado inicialmente por Helm et
 282 al. (1991).

283 No estudo de Beutin et al. (2007), foi possível demonstrar que o FT-IR pode
 284 revelar diferenças estruturais de superfície entre dois grupos de cepas de *E. coli*
 285 O123. Para Abu-Aqil et al. (2022) e Abu-Aqil et al. (2023), foi utilizado o FT-IR e
 286 o aprendizado de máquina para identificar *E. coli* diretamente da urina de
 287 pacientes hospitalizados, bem como determinar o perfil de suscetibilidade aos
 288 antibióticos e a presença de isolados produtores de beta-lactamases de espectro
 289 estendido.

290 O estudo de Helm et al. (1991) determinou cinco janelas espectrais
 291 correspondentes à absorção expressa em números de onda (cm⁻¹): janela 1,
 292 3000–2800cm⁻¹ dominada por vibrações presentes em ácidos graxos (lipídios);
 293 janela 2, 1800–1500cm⁻¹, dominada por vibrações das bandas amida I e amida II
 294 (proteínas e peptídeos); janela 3, 1500–1200cm⁻¹, com informações de
 295 proteínas, ácidos graxos e compostos de fosfato (região mista); janela 4, 1200–
 296 900cm⁻¹, bandas de carboidratos presentes na parede celular (polissacarídeos);
 297 janela 5, 900–700cm⁻¹, mostrando alguns padrões espectrais específicos, ainda
 298 não atribuídos a componentes celulares ou grupos funcionais (região de
 299 impressão digital).

300 Eses padrões específicos de impressão digital tornam este método uma
301 alternativa para identificação rápida de bactérias, tipagem e a triagem no nível
302 de subespécies, sendo as janelas 3 e 4, o ponto chave para a identificação
303 bacteriana (MEYERS, 2006; SHI et al., 2020).

304 Apesar da grande aplicabilidade de FT-IR em microbiologia nos últimos anos,
305 esta técnica pode ser considerada pouco convencional e seu potencial nas
306 rotinas de microbiologia pode ser subestimado, pela aceitação e adaptação
307 desta metodologia pela comunidade microbiológica. Entretanto, o método de FT-
308 IR poderia agregar valor à caixa de ferramentas da identificação bacteriana e,
309 em estudos futuros, nos processos de vigilância epidemiológica nacional
310 referentes à resistência bacteriana e a seus mecanismos (NOVAIS et al., 2019).

311

312 3 REFERÊNCIAS BIBLIOGRÁFICAS

- 313 ABRAHAM, E. P. et al. Further observations on penicillin. *Lancet*, 238, p. 177-
314 189, 1941.
- 315
- 316 ABU-AQIL, G. et al. Culture-independent susceptibility determination of *E.*
317 *coli* isolated directly from patients' urine using FTIR and machine-learning.
318 *Analyst*, v. 147, n. 21, p. 4815-4823. 2022.
- 319
- 320 ABU-AQIL, G. et al. Instant detection of extended-spectrum β-lactamase-
321 producing bacteria from the urine of patients using infrared spectroscopy
322 combined with machine learning. *Analyst*, v. 148, n.05, p. 1130-1140. 2023.
- 323
- 324 ADEYANKINNU, F. A. et al. A Multicenter Study of Beta-Lactamase Resistant
325 *Escherichia coli* and *Klebsiella pneumoniae* Reveals High Level Chromosome
326 Mediated Extended Spectrum β Lactamase Resistance in Ogun State, Nigeria.
327 *Interdisciplinary perspectives on infectious diseases*, v. 2014, 7p, 2014.
- 328
- 329 ANDRADE, L. N.; DARINI, A. L. C. Resistência bacteriana – Parte 1:
330 Mecanismos de Resistência aos antibióticos – Resistência bacteriana – Parte 2:
331 Conceitos e Definições. *Journal of Infection Control*, v. 7, n. 3, 2016.
- 332
- 333 BENGTSSON-PALME, J. et al. Using metagenomics to investigate human and
334 environmental resistomes, *Journal of Antimicrobial Chemotherapy*, v. 72, n. 10,
335 p. 2690–2703, 2017.
- 336
- 337 BEUTIN, L.; WANG, Q.; NAUMANN, D. et al. Relationship between O-antigen
338 subtypes, bacterial surface structures and O-antigen gene clusters in *Escherichia*
339 *coli* O123 strains carrying genes for Shiga toxins and intimin. *Journal of Medical*
340 *Microbiology*, v. 56, p.177–184, 2007.
- 341

- 342 BOONYASIRI, A. et al. Prevalence of antibiotic resistant bacteria in healthy
343 adults, foods, food animals, and the environment in selected areas in Thailand.
344 *Pathogens and Global Health*, v. 108, p.235–45, 2014.
- 345
- 346 BUNSEN, R.; KIRCHHOFF, G. R. Chemical analysis of substances by
347 observations on their spectra. *The Edinburgh Review or Critical Journal*, v. CXVI,
348 p.149-160, Jul – Oct. 1862.
- 349
- 350 BUSH, K.; JACOBY, G. A. Update functional classification of beta-
351 lactamases. *Antimicrobial Agents and Chemotherapy*, v. 54, n. 3, p. 969-76,
352 2010.
- 353
- 354 BLANCHARD, P.C. 2012. Diagnostics of dairy and beef cattle diarrhea.
355 *Veterinary Clinic of North America Food Animal Practice*, v.28, p443-464, 2012.
- 356
- 357 CANTÓN, R.; GONZÁLEZ-ALBA, J.M.; GALÁN, J.C. CTX-M Enzymes: Origin
358 and Diffusion. *Frontiers in Microbiology*, v.3, article 110, Apr. 2, 2012.
- 359
- 360 CARATTOLI, A. Animal reservoirs for extended spectrum beta-lactamase
361 producers. *Clinical Microbiology and Infection*, Jan; 14 Suppl 1:117-23, 2008.
- 362
- 363 COLOBATIU, L. et al. First description of plasmid-mediated quinolone resistance
364 determinants and beta-lactamase encoding genes in non-typhoidal *Salmonella*
365 isolated from humans, one companion animal and food in Romania. *Gut*
366 *Pathogens*, v. 16, n. 07. 2015.
- 367
- 368 CONSTABLE, P.D. Antimicrobial use in the treatment of calf diarrhea. *Journal of*
369 *Veterinary Internal Medicine*, v. 18, p. 8-17, 2004.
- 370
- 371 COQUE, T. M.; BAQUERO, F.; CANTON, R. Increasing prevalence of ESBL-
372 producing Enterobacteriaceae in Europa. *Eurosurveillance*, v. 13, n.47, 19044,
373 Nov. 20, 2008.
- 374
- 375 COURA, F. M. Longitudinal study of *Salmonella* spp., diarrheagenic *Escherichia*
376 *coli*, Rotavirus, and Coronavirus isolated from healthy and diarrheic calves in a
377 Brazilian dairy herd. *Tropical Animal Health Production*, v.47, p. 3-11, 2015.
- 378
- 379 CHONG, Y.; SHIMODA, S.; SHIMONO, N. Current epidemiology, genetic
380 evolution, and clinical impact of extended-spectrum β -lactamase-producing
381 *Escherichia coli* and *Klebsiella pneumoniae*. *Infection, Genetic and Evolution*,
382 v.61, p.185–8, 2018.
- 383
- 384 CLSI, CLINICAL AND LABORATORY STANDARDS INSTITUTE. Performance
385 Standards for Antimicrobial Susceptibility Testing. 31st ed. CLSI supplement
386 M100, v. 41, n. 03. USA, 2021.
- 387
- 388 EUCAST, European Committee on Antimicrobial Susceptibility Testing. *Expert*
389 *rules and expected phenotypes*. Disponível em:

- 390 https://www.eucast.org/expert_rules_and_expected_phenotypes. Acesso em: 02
391 março de 2022.
- 392
- 393 ELLER, C. et al. Emergence of extended-spectrum β -lactamase (ESBL) CTX-M-
394 8 in Germany. *Journal of Antimicrobial Chemotherapy*, v.69, n.2, p.562-4, Feb.
395 2014.
- 396
- 397 ELSHAFIEE, E. A. et al. carbapenem-resistant *Pseudomonas aeruginosa*
398 originating from farm animals and people in Egypt. *Journal Veterinary Research*,
399 v. 63, 2019.
- 400
- 401 EHRLICH, P. Address in Pathology, ON CHEMOTHERAPY: Delivered before
402 the Seventeenth International Congress of Medicine. *British Medical Journal*.
403 v.16, n.2, 1913.
- 404
- 405 EWERS, C. et al. OXA-23 and ISAb1–OXA-66 class D β -lactamases in
406 Acinetobacter baumannii isolates from companion animals. *International Journal*
407 of *Antimicrobial Agents*, v. 49, n. 1, p.-37-44, 2017.
- 408
- 409 FERNANDES, M. R. et al., Zooanthroponotic Transmission of Drug-Resistant
410 *Pseudomonas aeruginosa*, Brazil. *Emerging Infectious Diseases*. v, 24, n. 06, p.
411 1160-1162, 2018.
- 412
- 413 FLEMING, A. On the antibacterial action of cultures of a penicillium, with special
414 reference to their use in the isolation of *B. Influenzae*. *British Journal of*
415 *Experimental Pathology*, v.10, p. 226-236. 1929.
- 416
- 417 FRASAO, B. da SILVA et al. Detection of fluoroquinolone resistance by mutation
418 in *gyrA* gene of *Campylobacter*spp. Isolates from broiler and laying (*Gallus gallus*
419 *domesticus*) hens, from Rio de Janeiro State, Brazil. *Ciência Rural*, v. 45, n. 11,
420 p. 2013-2018, nov, 2015.
- 421
- 422 GUNDTRAN, R. S. et al. Prevalence and distribution of *bla*_{CTX-M}, *bla*_{SHV}, *bla*_{TEM}
423 genes in extended-spectrum β -Lactamase- producing *E. coli* isolates from broiler
424 farms in the Philippines. *BMC Veterinary Research*, v.15, n.227, p.1-8, 2019.
- 425
- 426 HAN, R. et al. Elimination kinetics of ceftiofur hydrochloride in milk after an 8-day
427 extended intramammary administration in healthy and infected cows. *PLoS One*,
428 v.12, n.11, e0187261, Nov. 2, 2017.
- 429
- 430 HELM, D.; LABISCHINSKI, H.; SCHALLEHN, G. et al. Classification and
431 identification of bacteria by Fourier-transform infrared spectroscopy. *Journal of*
432 *General Microbiology*, v. 137, p. 69-79, 1991.
- 433
- 434 IBGE/PPM, *Produção de Pecuária Municipal*. Rio de Janeiro, v. 48, p-1-12, 2020.
- 435
- 436 JARLIER, V. et al. Extended broad-spectrum betalactamases conferring
437 transferable resistance to newer beta-lactam agents in Enterobacteriaceae:
438 hospital prevalence and susceptibility patterns. *Reviews of Infectious Disease*, v.
439 10, p. 867–878. 1988.

- 440
441 KATZ, L.; BALTZ, R. H. Natural product discovery: past, present, and future.
442 *Journal of Industrial Microbiology and Biotechnology*, v. 43, p. 155-176, 2016.
443
444 LIVERMORE, D. M. et al. Interpretative reading: recognizing the unusual and
445 inferring resistance mechanisms from resistance phenotypes. *Journal of*
446 *Antimicrobial Chemotherapy*, v.1, p.87-102, 2001.
447
448 MAQUELIN K. et al. Identification of medically relevant microorganisms by
449 vibrational spectroscopy. *Journal of Microbiology Methods*, v.51, p. 255–271,
450 2002.
451
452 MEYERS, R. A. Encyclopedia of Analytical Chemistry (Applications, Theory, and
453 Instrumentation). *Infrared Spectroscopy in Microbiology.*, p. 1–32, 2006.
454
455 MELO, L. C. et al. Prevalence and molecular features of ESBL/pAmpC-producing
456 Enterobacteriaceae in healthy and disease companion animals in Brazil.
457 *Veterinary Microbiology*, v. 221, p. 59-66. 2018.
458
459 NEWTON, I. A Letter of Mr. Isaac Newton, Professor of the Mathematicks in the
460 University of Cambridge; Containing His New Theory about Light and Colors:
461 Sent by the Author to the Publisher from Cambridge, Febr. 6. 1671/72; In Order
462 to be Communicated to the R. Society. *Philosophical Transactions*, v. 6, p. 3075–
463 3087, 1972.
464
465 NOVAIS, A.; FREITAS, A. R.; RODRIGUES, C. et al. 27Fourier transform infrared
466 spectroscopy: unlocking fundamentals and prospects for bacterial strain typing.
467 *European Journal of Clinical Microbiology & Infection Diseases*, v.38, n.3, p.427-
468 448, 2019.
469
470 OLIVEIRA, C. J. B. et al. Methicillin-resistant *Staphylococcus aureus* from
471 Brazilian dairy farms and identification of novel sequence types. *Zoonoses Public*
472 *Health*, v. 63, p.97–105. 2016.
473
474 OTTEN, H. Domagk, and the development of the sulphonamides. *Journal*
475 *Antimicrobial Chemotherapy*, v.17, p. 689-690, 1986.
476
477 O'NEILL, J. *Tackling drug-resistant infections globally: final report and*
478 *recommendations. Review on Antimicrobial Resistance*. Wellcome Trust and HM
479 Government. Disponível em:
480 https://amrreview.org/sites/default/files/160525_Final%20paper_with%20cover.pdf. 2016.
481
482
483 PALMEIRA, J.D., et al. Epidemic spread of IncI1/Pst113 plasmid carrying the
484 Extended-Spectrum Beta-Lactamase (ESBL) *bla*CTX-M-8 gene in *Escherichia*
485 *coli* of Brazilian catte. *Veterinary Microbiology*, v. 243, 108629, April 2020.
486
487 PATERSON, D. L.; BONOMO, R. A. Extended-spectrum beta-lactamases: a
488 clinical update. *Clinical Microbiology Reviews*, v.8, n.4, p.657-86, 2005.

- 489
490 PITOUT, J.D.; CHURCH, D. L.; GREGSON, D. B. et al. Molecular epidemiology
491 of CTX-M-producing *Escherichia coli* in the Calgary Health Region: emergence
492 of CTX-M-15-producing isolates. *Antimicrobial Agents Chemotherapy*, v. 51, n. 4,
493 p.1281-6, 2007.
- 494
495 POMBA, C. et al. First Report of OXA-23-Mediated Carbapenem Resistance in
496 Sequence Type 2 Multidrug-Resistant *Acinetobacter baumannii* Associated with
497 Urinary Tract Infection in a Cat. *Antimicrobial Agents and Chemotherapy*, vol. 58,
498 n. 2, 2015.
- 499
500 POIREL, L. et al. Antimicrobial resistance in *Escherichia coli*. *Microbiology
501 Spectrum Journal*, v.6, n.4, ARBA-0026-2017. 2018.
- 502
503 PRIBUL, B. R. et al. Characteristics of quinolone resistance in *Salmonella* spp.
504 Isolates from the food chain in Brazil. *Frontiers in Microbiology*, v. 8, p. 299. Mar
505 14, 2017.
- 506
507 ROCHA, F.R.; PINTO, V.P.; BARBOSA, F.C. The Spread of CTX-M-Type
508 Extended-Spectrum β-Lactamases in Brazil: A Systematic Review. *Microbial
509 Drug Resistance*, v.22, n.4, p.301-11, Jun. 2016.
- 510
511 SANTAJIT, S.; INDRAWATTANA, N. Mechanisms of Antimicrobial Resistance in
512 ESKAPE pathogens. *BioMed research international*, v. 2016, p.1-9, 2016.
- 513
514 SAUGET, M. et al. A espectrometria de massa MALDI-TOF pode razoavelmente
515 classificar bactérias? *Trends in Microbiology*, v. 25, p.447–455, 2017.
- 516
517 SABAT, A. J.; BUDIMIR, A.; NASHEV, D. et al. Overview of molecular typing
518 methods for outbreak detection and epidemiological surveillance.
519 *Eurosurveillance*, v. 18, n. 4, 2013.
- 520
521 SUKMAWINATA, E. et al. Incl1 Plasmid Associated with *bla*_{CTX-M-2} Transmission
522 in ESBL-Producing *Escherichia coli* Isolated from Healthy Thoroughbred
523 Racehorse, Japan. *Antibiotics*, v.9, n.70, p. 2-7, 2020.
- 524
525 SMITH T.; ORCUTT M. L. The bacteriology of the intestinal tract of young calves
526 with special reference to the early diarrhea ("scours"). *Journal of Experimental
527 Medicine*, v. 41, p.89–106, 1925.
- 528
529 SCHMID, A.; HÖRMANSDORFER, S.; MESSELHÄUSSER, U. Prevalence of
530 extended-spectrum β-lactamase-producing *Escherichia coli* on Bavarian dairy
531 and beef cattle farms. *Applied and Environmental Microbiology*, v.79, n.9, p.3027-
532 32, 2013.
- 533
534 SHI, H. et al. The strategy for correcting interference from water in Fourier
535 transform infrared spectrum based bacterial typing. *Talanta*, v. 208, 2020.
- 536

- 537 STUART, B. *Infrared Spectroscopy: Fundamentals and Applications*. John Wiley
538 & Sons. West Sussex. England. 2004.
- 539
- 540 SETTLE F. A. *Handobook of Instrumental Techiniques for Analytical Chemistry*.
541 Prentice Hall. New Jersey. USA.1997.
- 542
- 543 SPINOSA, H. de SOUZA.; GÓRNIAK, S. L.; BERNARDI, M. M. *Farmacologia*
544 *Aplicada à Medicina Veterinária*. 5^a Ed. Rio de Janeiro: Guanabara Koogan,
545 2014.
- 546
- 547 TEKINER, L.; OZPINAR, H. Occurrence and characteristics of extended
548 spectrum beta-lactamases-producing Enterobacteriaceae from foods of animal
549 origin. *Brazilian Journal of Microbiology*, v. 47, p.444-51, 2016.
- 550
- 551 TUTIJA, J. F et al. Molecular and phenotypic characterization of *Escherichia coli*
552 from calves in an important meat-producing region in Brazil. *The Journal of*
553 *Infection in Developing Countries*, v.16, n.6, p.1030-1036. Jun 30, 2022.
- 554
- 555 WIDODO, A.; EFFENDI, M. H.; KHAIRULLAH, A.R. Extended-spectrum beta-
556 lactamase (ESBL)-producing *Eschericia coli* from livestock. *Systematic Reviews*
557 *in Pharmacy*, v. 11, n. 7, p. 382-392, 2020.
- 558
- 559 WHO, WORLD HEALTH ORGANIZATION. *The envolving threat of antimicrobial*
560 *resistance options for action*. 2012. Disponível em:
561 http://apps.who.int/iris/bitstream/handle/10665/44812/9789241503181_eng.pdf;jsessionid=48D7D6DA4B598117F22522019F328FB7?sequence=1. Acesso em:
562 27 de fevereiro de 2022.
- 563
- 564
- 565 WHO, WORLD HEALTH ORGANIZATION. *Global Action Plan on Antimicrobial*
566 *Resistance*. Geneva, Switzerland. World Health Organization, 2017.

567

CAPÍTULO 2

568

4. (ARTIGO I)

569

571 [Artigo formatado de acordo com as normas da Revista Arquivo Brasileiro de Medicina
572 Veterinária e Zootecnia]

573

IDENTIFICAÇÃO FENOTÍPICA E GENOTÍPICA DE *Escherichia coli* PRODUTORAS DE BETA-LACTAMASES DE ESPECTRO ESTENDIDO (ESBL) ISOLADAS DE FEZES DE BEZERROS

577

578 *Phenotypic and genotypic characterization of Escherichia coli producing extended
579 spectrum beta-lactamases (ESBL) isolated from calf feces.*

580

581 MARANGONI, Y. G.*¹, RAMOS, C.A.N.³, TUTIJA, J. F.¹, BENITES, C. E. S.², LEAL,
582 C. R. B.*³

583

584 ¹Programa de Pós-Graduação em Ciências Veterinárias - Universidade Federal de Mato
585 Grosso do Sul – Campo Grande, MS, Brasil.

586 ²Aluno de graduação - Universidade Federal de Mato Grosso do Sul – Campo Grande,
587 MS, Brasil.

588 ³Docentes da Universidade Federal de Mato Grosso do Sul – Campo Grande, MS, Brasil.

589 *yasmin.marangoni@ufms.br ; *cassia.leal@ufms.br

590

591

RESUMO

592 A produção de beta-lactamases por Enterobacteriales é o mecanismo de resistência mais
593 comum aos antibióticos beta-lactâmicos. As beta-lactamases de espectro estendido
594 (ESBL), são consideradas uma urgência clínica e epidemiológica mundial. Este estudo
595 teve como objetivo verificar a presença de cepas de *Escherichia coli* diarreogênicas,
596 isoladas de bezerros, produtoras de ESBL. Amostras de fezes de 208 bezerros, de duas
597 faixas etárias distintas, foram obtidas e cultivadas em meio de enriquecimento para
598 isolamento de *E. coli*. A análise fenotípica foi realizada pelo método de triagem de disco
599 de aproximação empregando discos de amoxicilina + ácido clavulônico (AMC),
600 ceftriaxona (CRO), cefotaxime (CTX), ceftazidima (CAZ) e aztreonam (ATM). Foi
601 realizada a Reação em Cadeia da Polimerase (PCR), para identificação dos genes de
602 resistência *bla*CTX-M-2, *bla*CTX-M-8, *bla*CTX-M-15, *bla*SHV e *bla*TEM. A presença de pelo menos
603 um dos genes citados foi verificada em 35,6% do total (n=74). A distribuição dos genes
604 nas amostras positivas ocorreu da seguinte forma: 3,84% (n=8) para *bla*CTX-M-2, 5,28%
605 (n=11) para *bla*SHV e 29,3% (n=61) para o gene *bla*TEM. Os genes *bla*CTX-M-15 e *bla*CTX-M-8

606 não foram observados no estudo. Quanto à suscetibilidade aos antibióticos, as amostras
607 apresentaram altas taxas de resistência ao grupo dos beta-lactâmicos, bem como para
608 tetraciclina (80%). Foi possível verificar que 50,9% (106/208) apresentaram perfil de
609 multirresistência. Este estudo demonstra que bezerros podem ser reservatórios de cepas
610 de *E. coli* produtoras de ESBL, constituindo um risco à saúde humana e animal, pelo risco
611 de disseminação desses isolados no ambiente.

612 **Palavras-chave:** beta-lactâmicos, ESBL, pecuária e resistência à antibióticos.

613

614 ABSTRACT

615

616 Beta-lactamase production by Enterobacteriales is the most common resistance
617 mechanism for beta-lactam antibiotics. Extended spectrum beta-lactamase (ESBL) are
618 considered a world clinical and epidemiological urgency. This study aimed to verify the
619 presence of diarrheagenic *Escherichia coli* strains, isolated from calves, ESBL producers.
620 Fecal samples from 208 calves, from two distinct age groups, were obtained and
621 cultivated in enrichment for *E. coli* isolation. Phenotypic analysis was performed by the
622 approximation disc screening method employing amoxicillin discs + clavulonic acid
623 (AMC), ceftriaxone (CRO), cefotaxime (CTX), ceftazidima (CAZ) and aztreonam
624 (ATM). Polymerase Chain Reaction (PCR) was performed to identify the *bla*_{CTX-M-2},
625 *bla*_{CTX-M-8}, *bla*_{CTX-M-15}, *blas*_{HV} and *bla*_{TEM} resistance genes. The presence of at least one of
626 the genes cited was verified in 35.6% of the total (n = 74). The distribution of genes in
627 the positive samples occurred as follows: 3.84% (n = 8) to *bla*_{CTX-M-2}, 5.28% (n = 11) for
628 *blas*_{HV} and 29.3% (n = 61) for *bla*_{TEM} gene. The *bla*_{CTX-M-15} and *bla*_{CTX-M-8} genes were not
629 observed in the study. Regarding susceptibility to antibiotics, the samples showed high
630 resistance rates to the beta-lactam group, as well as for tetracycline (80%). It was possible
631 to verify that 50.9% (106/208) had multipurpose profile. This study demonstrates that
632 calves can be *E. coli* strains reservoirs of ESBL producing, constituting a risk to human
633 and animal health, due to the risk of dissemination of these isolates in the environment.

634 **Keywords:** beta-lactams, ESBL, livestock and antibiotic resistance.

635

636 INTRODUCTION

637

638 The identification of multiresistant bacteria and their virulence genes in livestock
639 has great importance in public health worldwide, as well as the identification of possible

640 antibiotic resistance genes, resulting from exacerbated therapeutic conduct and selective
641 pressure (Han *et al.*, 2017).

642 Diarrhea in calves is one of the main causes of economic losses in Brazilian and
643 worldwide livestock. Since the beginning of the 20th century, the worldwide scientific
644 community has pointed to the bacterium *Escherichia coli* as one of the main agents
645 involved in diarrhea of infectious origin, therefore, this bacterial genus and its
646 diarrheagenic strains have been studied and identified (Smith; Orcutt, 1925; Coura *et al.*,
647 2015; Tutija *et al.*, 2020).

648 The differentiation between *E. coli* pathogenic strains from non-pathogenic strains
649 is based on the production of virulence factors, as well as on the identification of the
650 mechanisms related to disease. In response to the widespread use of antimicrobials in the
651 treatment of enteritis cases in calves, spread of virulence factors between herds may
652 occur. This event may be relevant for the emergence and dissemination of bacteria
653 capable of expressing enzymes that confer resistance to multiple antibiotics (Coura *et al.*,
654 2015; Poirel *et al.*, 2018).

655 Several studies have pointed to the worldwide growth of this resistance by
656 members of Enterobacteriales, such as the emergence of ESBL, which can irreversibly
657 hydrolyze the beta-lactam ring of the amine bond, preventing its action on transpeptidases
658 (Bush and Jacoby, 2010). They can hydrolyze all penicillins, cephalosporins including
659 third and fourth generation and aztreonam (Coque *et al.*, 2008).

660 Feces of production animals, which have strains with antimicrobial resistance
661 genes, such as the ESBL enzyme, can be constant reservoirs of dissemination to humans
662 (Carattoli, 2008; Schmidt *et al.*, 2013), the environment and other animals (Boonyasiri *et*
663 *al.*, 2014; Tekiner; Ozpinar, 2016; Melo *et al.*, 2018; Brisola *et al.*, 2019), leading to
664 delays in appropriate therapy, increased morbidity, mortality, and consequently,
665 economic losses to producers and industry (Santajit; Indrawattana, 2016; Chong *et al.*,
666 2018).

667 Thus, the aim of this study was to evaluate the presence of diarrheagenic *E. coli*
668 strains producing ESBL, from an important beef cattle producing region in Brazil.

669

670 MATERIAL AND METHODS

671

672 Two hundred and eight isolates ($n = 208$) of diarrheagenic *E. coli* were used in
673 this study. These isolates were previously characterized as diarrheagenic *E. coli* by Tutija
674 *et al.* (2020).

675 These isolates came from stool and rectal swabs, from beef calves, with ages
676 varying from one to 60 days. The study was conducted at the Veterinary Bacteriology
677 Laboratory at the Universidade Federal de Mato Grosso do Sul (UFMS), from March
678 2021 to December 2022.

679 In vitro susceptibility to antimicrobials was verified, according to the antibiogram
680 method standardized by the CSLI (Clinical Laboratory Standard Institute) (2021), which
681 is based on the method originally described by Bauer *et al.* (1966). The most used
682 antibiotics in the field reported by the veterinarians themselves were used, such as:
683 amoxicillin + clavulanic acid (AMC), amoxicillin (AMO), cephalexin (CFE),
684 enrofloxacin (ENO), florfenicol (FLF), gentamicin (GEN), norfloxacin (NOR), penicillin
685 G (PEN), sulfamethaxazol+trimethoprim (SUT) and tetracycline (TET).

686 As for the phenotypic analysis of ESBL-encoding genes, the approximation disk
687 screening technique according to Jarlier *et al.* (1988), was performed using amoxicillin +
688 clavulanic acid (AMC), ceftriaxone (CRO), cefotaxime (CTX), ceftazidine (CAZ) and
689 aztreonam (ATM) discs. A sample of *E. coli* ATCC® 25922 was used as a control,
690 verifying the interaction of the AMC antibiotic with the others.

691 Subsequently, molecular analysis of the diarrheagenic *E. coli* isolates was
692 performed to identify the genes encoding the ESBL enzyme, such as *bla*_{CTX-M} (including
693 *bla*_{CTX-M-2}, *bla*_{CTX-M-8} and *bla*_{CTX-M-15}), *blas*_{HV} and *bla*_{TEM}, using Reaction Polymerase in
694 Chain (PCR).

695 The isolates were suspended in BHI broth (brain and heart infusion), incubated in
696 a culture oven for 24 hours. After this time, the centrifugation was performed and the
697 extraction of the sediment DNA started, according to the protocol described by Adwan
698 (2014). For the analysis of DNA purity and its quantification, the NanoDrop™ OneC
699 (Thermo Fischer Scientific) was used.

700 For the amplification of target genes (Tab. 1), the reactions were performed with
701 a final volume of 25 μ L, containing 2.5 μ L of 10X buffer (20 mM Tris-HCl, pH 8.3, 50
702 mM KCl), 1.5 mM MgCl₂, 0.2 mM dNTP, 1.25 U of *Taq* DNA polymerase (5 U/uL), 10
703 pmol of each primer (100 ng/uL) and 2 μ L of DNA (approximately 50 ng).

704 Amplification conditions for the CTX-M-like enzyme included: 1) initial
705 denaturation at 94°C for 5 minutes; 2) denaturation at 94°C for 1 minute; 3) annealing at

706 a specific temperature for each primer, for 1 minute; 4) extension at 72°C for 1 minute;
 707 5) repetitions of steps 2, 3 and 4, 29 times; ending with step 6, the final extension at 72°C,
 708 for 5 minutes.

709 As for the SHV-type enzyme, they included: 1) initial denaturation at 94°C for 5
 710 minutes; 2) denaturation at 94°C for 30 seconds; 3) annealing at a specific temperature
 711 for each initiator, for 30 seconds; 4) extension at 72°C for 1 minute; 5) repetitions of steps
 712 2, 3 and 4, 32 times; with final extension at 72°C for 10 minutes.

713 For the TEM-type enzyme, they included: 1) initial denaturation at 94°C for 5
 714 minutes; 2) denaturation at 95°C for 30 seconds; 3) annealing at a specific temperature
 715 for each primer, for 1 minute; 4) extension at 72°C for 2 minutes; 5) repetitions of steps
 716 2, 3 and 4, 32 times; with final extension at 72°C for 5 minutes.

717

718 **Table 1. Initiators to identify the genes encoding ESBLs.**

| Genes | Sequence 5'-3' | T (°C) * | bp** | References |
|--------------------------------|--|----------|------|--------------------------------------|
| <i>bla</i> _{CTX-M-2} | F5'-GCGACCTGGTTAACTACAATC-3' R5'-CGGTAGTATTGCCCTTAAGCC-3' | 55 | 351 | Torrentino <i>et al.</i> , 2011 |
| <i>bla</i> _{CTX-M-8} | F5'-CTGGAGAAAAGCAGCGGGGG-3' R5'-ACCCAGGATGTGGGTAGCCC-3' | 55 | 320 | Bonnet <i>et al.</i> , 2000 |
| <i>bla</i> _{CTX-M-15} | F5'-CACACGTGGAATTAGGGACT-3' R5'-GCCGTCTAAGGCGATAAACAA-3' | 56 | 550 | Muzahed <i>et al.</i> , 2008 |
| <i>bla</i> _{SHV} | F5'-AGGATTGACTGCCTTTTG-3' R5'-ATTGCTGATTCGCTCG3' | 54 | 392 | Colom <i>et al.</i> , 2003 |
| <i>bla</i> _{TEM} | F5'-TTGGGTGCACGAGTGGGTTA-3' R5'-TAATTGTTGCCGGGAAGCTA-3' | 54 | 465 | Gangoue-Pieboji <i>et al.</i> , 2005 |

719 *T (°C): annealing temperature; ** bp: base pairs.

720

721 Subsequently, the amplified products were analyzed by electrophoresis in agarose
 722 gel 1%, together with *E. coli* ATCC® 25922. Molecular weight marker (Ladder 100bp,
 723 Ludwing Biotec, Brazil), and Gel Red nucleic acid stain (Biotium, USA) were used. The
 724 amplified products were visualized under ultraviolet light, with the aid of a transilluminator
 725 (Transiluminator UV for gel - LTB HE, Loccus, Brazil).


726

727 RESULTS

728

729 In the phenotypic analysis, an increase in the diameter of the inhibition zone
730 towards the AMC disk or the appearance of the phantom zone was observed in 31.25%
731 (65/208) samples, as shown in the figure 1.

732




733

734 Figure 1: Positive approximation disk screening technique demonstrating the phantom zone or absence of
735 inhibition halo (red arrow) to verify samples possibly producing the ESBL enzyme.

736

737 Genotypic analysis by the PCR method for the search for ESBL coding enzymes
738 showed positivity in 3.84% (8/208) of the samples for the *blaCTX-M-2* enzymes, 5.28%
739 (11/208) for the gene *blaSHV* and 29.3% (61/208) for the *blaTEM* gene (Figure 2). The
740 *blaCTX-M-15* and *blaCTX-M-8* genes were not observed in the study.

741




742

743 Figure 2: Electrophoresis of positive samples on a 1% agarose gel, using a molecular marker of 100 base
744 pairs, visualizing the amplification of the *blaCTX-M-2* and *blaSHV* (A), as well as the *blaTEM* gene (B).

745

746 Through genetic identification, it was possible to observe that in the four macro
747 regions of the state of Mato Grosso do Sul, at least one ESBL coding gene was located
748 (Figure 3).

749



750

751 Figure 3: Map of the state of Mato Grosso do Sul, Brazil, showing the positive municipalities (yellow) for
 752 the genotypic identification of the ESBL-encoding genes: *blaCTX-M-2* (triangle), *blaSHV* (square) and *blaTEM*
 753 (circle).

754

755 In this study, it was possible to observe that 65.7% of the positive samples for
 756 ESBL coding genes are from calves between 1 and 30 days of age and 34.2% between 31
 757 and 60 days.

758 In 35.6% (74/208) of the samples positive for the enzymatic variants CTX-M,
 759 SHV and TEM, the frequency of in vitro antimicrobial resistance of the class of beta-
 760 lactams tested can be observed (Tab. 2A) and, additionally, the correlation of the
 761 frequency of in vitro antimicrobial resistance of antimicrobials of the aminoglycosides,
 762 fluoroquinolones, tetracyclines, amphenicols and sulfonamides classes, with the
 763 localization of CTX-M, SHV and TEM family genes (Tab. 2B).

764

765 Table 2. Correlation of the frequency of antimicrobial resistance to beta-lactams (A) by the disk diffusion
 766 method and antimicrobials from the classes of aminoglycosides, fluoroquinolones, tetracyclines,
 767 amphenicols and sulfonamides (B), with the location of the *blaCTX-M-2* genes, *blaSHV* and *blaTEM* from
 768 diarrheagenic *E. coli* isolates by the PCR method.

A

| Antibiotics | Number of resistant samples (%) | | |
|--------------|---------------------------------|---------------|---------------|
| | <i>blaCTX-M-2</i> | <i>blaSHV</i> | <i>blaTEM</i> |
| Penicillin G | 8 (100%) | 11 (100%) | 61 (100%) |
| Amoxicillin | 7 (87.5%) | 4 (36.36%) | 43 (70.49%) |

| | | | |
|-------------------------------|-----------|------------|-------------|
| Cephalexin | 7 (87.5%) | 6 (54.54%) | 23 (37.70%) |
| Amoxicillin + Clavulanic Acid | 3 (37.5%) | 0 | 16 (26.22%) |
| B | | | |
| Tetracycline | 7 (87.5%) | 7 (63.63%) | 55 (90.16%) |
| Sulfamethoxazole+Trimethoprim | 4 (50%) | 4 (36.36%) | 35 (57.37%) |
| Enrofloxacin | 4 (50%) | 6 (54.54%) | 22 (36.06%) |
| Norfloxacin | 3 (37.5%) | 5 (45.45%) | 17 (27.86%) |
| Gentamicin | 3 (37.5%) | 5 (45.45%) | 14 (22.95%) |
| Florfenicol | 2 (25%) | 0 | 7 (11.47%) |

769

770 After performing the antibiogram, the frequency of in vitro antimicrobial
 771 resistance of the 208 samples was observed (Tab. 3), in which the highest resistance rate
 772 was for penicillin G (100%), followed by tetracycline (80%), amoxicilin (61%),
 773 sulfametaxazole+trimethoprim (45.7%) and cephalexin (45%) and with less frequency of
 774 antimicrobial resistance, florfenicol (12%).

775

776 Table 3. Frequency of in vitro antimicrobial resistance in 208 diarrheagenic *Escherichia coli* isolates from
 777 stool and rectal swab samples.

| Antibiotics | Number of resistant samples (%) |
|-------------------------------|---------------------------------|
| Penicilin | 208 (100%) |
| Tetracycline | 166 (80%) |
| Amoxicillin | 126 (61%) |
| Cephalexin | 94 (45%) |
| Sulfamethoxazole+Trimethoprim | 95 (45.7%) |
| Enrofloxacin | 66 (32%) |
| Norfloxacin | 50 (24%) |
| Amoxicillin + Clavulanic Acid | 43 (20.6%) |
| Gentamicin | 39 (19%) |
| Florfenicol | 25 (12%) |

778

779 One hundred and six (50.9%) samples showed multidrug-resistance to three or
 780 more classes of antibiotics, in which no antibiotic was 100% effective, of these, 48.11%
 781 were positive for the phenotypic disk approximation test, and 42.45% molecularly
 782 positive for ESBL enzymes in relation to the presence of multidrug-resistance.

783 Twenty-eight (13.5%) samples of diarrheagenic *E. coli* were positive for the disk-
 784 approximation screening test and for the presence of one of the genes encoding ESBL.
 785 Forty-five samples (21.6%) did not express the phantom zone or lack of inhibition zone
 786 and were molecularly positive. However, 32 (15.4%) samples tested positive for the
 787 screening test and were molecularly negative.

788

DISCUSSION

790

791 Production animals, such as cattle, are considered one of the main reservoirs of *E.*
792 *coli* producing ESBL (Chong *et al.*, 2018).

793 According to the age of the animals, Brunton *et al.* (2014), demonstrated in their
794 study that resistance genes may be less prevalent in relation to the growth of the animals,
795 due to the increase in the physical area in which they remain, hindering the transmission
796 of resistance genes between them. In this case, the animals in the study were not weaned
797 and were in the same physical area, according to the selected farm. The limitation of the
798 age range in the present study, up to 60-days-old, may have influenced the non-alteration
799 of the indices. When extending this age range to more than 60 days, it is possible that the
800 variation mentioned by Brunton *et al.* (2014) could be noticed.

801 The situation of ESBL incidence in the world has undergone several changes, and
802 in the new epidemiological scenario, the prevalence of enzymes of the CTX-M family
803 stands out, both in hospital isolates, the community, and animals. This worldwide spread
804 was described as “the CTX-M pandemic” (Cantón *et al.*, 2012).

805 In the present study, enzymes of the CTX-M family were highlighted in 8 (3.84%)
806 of the 208 fecal samples from calves. The study by Sukmawinata *et al.* (2020) confirmed
807 the presence of this family in the feces of thoroughbred horses in Japan, as well as the
808 study by Gundran *et al.* (2019) which obtained 89.86% gene positivity in isolates from
809 broiler chickens in the Philippines, confirming the cosmopolitanization of the production
810 of ESBL type CTX-M.

811 In the molecular analysis, the prevalence of genes encoding ESBL was higher for
812 the *blaTEM* gene (29.3%), followed by *blasHV* (5.28%) and, lastly, by the *blaCTX-M-2* gene
813 (3.84%). The *blaCTX-M-15* and *blaCTX-M-8* genes were not observed in this study.

814 In Brazilian cattle, the *blaCTX-M-15* and *blaCTX-M-2* genes were found in the Southeast
815 region, mainly in the states of São Paulo and Rio de Janeiro (Rocha *et al.*, 2016), as well
816 as in the Northeast region (Palmeira *et al.*, 2020). Sukmawinata *et al.* (2020) reported a
817 high prevalence of the *blaCTX-M-2* gene, however, among the few studies that investigated
818 the prevalence of ESBL in calves, Eller *et al.* (2014) reported a high prevalence of *blaCTX-*
819 *M8*, unlike our study.

820 The studies reported in human and veterinary medicine by Cergole-Novella *et al.*
821 (2010), Rocha *et al.* (2016), Widodo *et al.* (2020), and Ejaz *et al.* (2021) demonstrated
822 that the prevalence of the *blaCTX-M*, *blaTEM* and *blasHV* gene is increasing among ESBL-
823 producing bacteria, that corroborates our study. They are common in humans (Aworh *et*
824 *al.*, 2022), especially in nosocomial infections; in animals - dogs, cats, horses, chickens,

825 pigs, cattle, wild animals (Ewers *et al.*, 2010; Melo *et al.*, 2018; Brisola *et al.*, 2019) - and
826 in the environment (Cantón *et al.*, 2012).

827 For Aworh *et al.* (2022), healthcare professionals should be aware that people who
828 have close contact with calves and their feces are a high-risk group for spreading *E. coli*
829 with ESBL, when compared to the general population. In their study, in addition to
830 isolating *E. coli* with ESBL in 45.4% of calves' feces from an abattoir, they observed the
831 presence of this enzyme in workers (41.2%) and in the work environment (13.4%).

832 The WHO (2017) lists bacterial agents and their respective antimicrobial
833 resistance in human medicine, but within veterinary medicine we can already observe the
834 cosmopolitanization of *Acinetobacter baumannii* (Pomba *et al.*, 2015; Ewers *et al.*, 2017),
835 *Pseudomonas aeruginosa* (Fernandes *et al.*, 2018; Elshafiee *et al.*, 2019), as well as
836 ESBL-producing *E. coli* (Melo *et al.*, 2018; Brisola *et al.*, 2019; Palmeira *et al.*, 2020) in
837 the critical priority group, and in the high priority group, *Staphylococcus aureus* (Oliveira
838 *et al.*, 2016), *Salmonella* species (Colobatiu *et al.*, 2015; Pribul *et al.*, 2017) and
839 *Campylobacter* species (Frasao *et al.*, 2015), at the human-animal-environment interface.

840 The production of ESBL by *E. coli* bacteria is a major public health problem, as
841 infections by these bacteria can result in failure of beta-lactam therapy (Chong *et al.*,
842 2018). Another reason is the phenomenon of co-resistance to other antibiotics, as they are
843 encoded by plasmids. Profiles of resistance genes for aminoglycoside, macrolides,
844 tetracyclines and fluoroquinolones antibiotics were observed by Carey *et al.* (2022) for
845 the *blactX-M* gene, which may facilitate co-selection processes, limiting current
846 therapeutic options (Paterson; Bonomo, 2005; Cantón *et al.*, 2012; Adeyankinnu *et al.*,
847 2014).

848 According to a report by the OIE (2021), tetracyclines were the most used
849 antibiotics in food-producing animals, followed by macrolides and penicillins.

850 Regarding antimicrobial resistance to the beta-lactam class and the location of the
851 genes that encode ESBL, we could observe that the highest index of resistance was for
852 penicillin G (100%), similar results to those of Shahrani *et al.* (2014), who evaluated the
853 antimicrobial resistance profile correlated with the identification of the STEC virulence
854 gene in samples of diarrhea from calves. However, it is known that members of the
855 Enterobacterales group, such as *E. coli*, are intrinsically resistant to penicillin G, that is,
856 this drug was being administered incorrectly, possibly due to the lack of knowledge of
857 the bacteria's intrinsic resistance.

Brower *et al.* (2017), Rahmahani *et al.* (2020), and Jarrige *et al.* (2020), observed a profile of significant resistance to tetracycline in poultry farms (47%), domestic chickens (48%), from cloacal swab, and in milk dairy calves (68%), results that corroborate the findings of this study, presenting antimicrobial resistance in 80% (166/208) for tetracycline. In addition, records from the farms tested showed that, at some point during management, tetracycline was administered to the calves in the study.

Bacteria showed the lowest frequency of resistance to florfenicol, with only 12% (25/208) of inhibition failure in the tested samples, as well as the study by Nespolo *et al.* (2014), who analyzed 52 bovine isolates destined for slaughter, obtained a resistance index of 1.92% to florfenicol.

E. coli is also prone to developing broad drug resistance, as it has a great capacity to accumulate resistance genes, mainly by horizontal gene transfer, which leads to infections that are difficult to treat (Poirel *et al.*, 2018). In the present study it was observed three samples with two tested genes, *blasHV* and *blaTEM*, and three samples with *blaCTX-M-2* and *blaTEM*, as well as their multidrug resistance to six classes of antibiotics, such as beta-lactams, fluoroquinolones, tetracycline, aminoglycoside, sulfonamides and amphenocol.

The multidrug-resistance profile, characterized by its resistance to three or more classes of antimicrobials (Magiorakos *et al.*, 2011), was found in 50.9% (106/208) of the isolates, results like Cergole-Novella *et al.* (2010) and Brisola *et al.* (2019), in which multidrug resistance was observed in eight of the 12 samples (66.7%) and 50 of the 135 samples (37.4%), respectively.

Al-Tamimi *et al.* (2019) proposes to use the phenotypic synergy test to detect the synergistic ability of clavulanic acid with cephalosporins and monobactam in ESBL-positive isolates. However, in our study there were divergences between positive samples in the phenotypic test (15.4%) and negative in the genotypic analysis, which could explain this phenomenon is the use of specific primers for the CTX-M, SHV and TEM families in this study, and the literature mentions other genes that can encode ESBL enzymes such as *blaCMY* (Brisola *et al.*, 2019) and *blaOXA* (Palmeira *et al.*, 2020).

In the phenotypic synergism test, the ideal distance for high synergistic activity is 20 to 25 mm between cephalosporins/monobactam and amoxicillin and clavulanic acid (Jarlier *et al.*, 1988; Bengtsson-Palme *et al.*, 2017). In the present study, 45 (21.6%) samples did not express the phantom zone or any zone of inhibition and were molecularly

891 positive, possibly greater distances could decrease the sensitivity of the test, hiding most
892 of the resistance factors during the laboratory routine (Al-Tamimi *et al.*, 2019).

893 Many production animals, carrying these strains of ESBL, can be constant
894 reservoirs of dissemination for humans, such as, for example, the contamination of
895 products of animal origin (Tekiner; Ozpinar, 2016).

896 Currently, there are no national programs for monitoring bacterial resistance and
897 its mechanisms in livestock, which makes it difficult to estimate the proportion of ESBL-
898 producing strains in the federations. In turn, human and animal infections by ESBL-
899 producing bacteria is associated with increased mortality, morbidity, high cost of
900 hospitalization, delay in adequate therapy and, consequently, economic losses for
901 producers (Santajit; Indrawattana, 2016; Chong *et al.*, 2018).

902

903 CONCLUSION

904

905 This study presents the first report on the identification of the *bla*_{CTX-M-2}, *bla*_{SHV}
906 and *bla*_{TEM} genes in stool samples from calves from Mato Grosso do Sul, Brazil, alerting
907 veterinarians about the use and knowledge of antibiotics in livestock, indicating the need
908 to establish strategies for surveillance and tracking of antibiotic resistance in the country.

909

910 ACKNOWLEDGMENTS

911

912 The present study was carried out with the support of the Universidade Federal de
913 Mato Grosso do Sul – UFMS/MEC, Brazil, Coordenação de Aperfeiçoamento de Pessoal
914 de Nível Superior (CAPES), code 001. Conselho Nacional de Desenvolvimento
915 Científico e Tecnológico (CNPQ), code 403651/2020-5; 302525/2022-0; 440214/2021-
916 1. Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia do Estado
917 de Mato Grosso do Sul (FUNDECT), code 007/2019; 360/2022.

918

919 REFERENCES

920

921 ADEYANKINNU, F. A. et al. A Multicenter Study of Beta-Lactamase Resistant
922 *Escherichia coli* and *Klebsiella pneumoniae* Reveals High Level Chromosome Mediated
923 Extended Spectrum β Lactamase Resistance in Ogun State, Nigeria. *Interdisciplinary*
924 *perspectives on infectious diseases*, v. 2014, 7p, 2014.

- 925
926 ADWAN, K. Fast DNA isolation and PCR protocols for detection of methicillin-resistant
927 staphylococci. *Folia Microbiology*, v.59, n.5-8, 2014.
- 928
929 AL-TAMIMI, M. et al. Effects of antimicrobial exposure on the antimicrobial resistance
930 of *Escherichia coli* in the digestive flora of dairy calves. *Microbial Drug Resistance*, v.
931 00, n. 00, 2019.
- 932
933 AWORH, M.K. et al. Extended-Spectrum β -Lactamase- Producing *Escherichia coli*
934 Among Humans, Beef Cattle, and Abattoir Environments in Nigeria. *Frontiers Cellular*
935 and Infection Microbiology, v. 12:869314. 2022.
- 936
937 BAUER A. W.; KIRBY W. M. M., SHERRIS. Antibiotic Susceptibility Testing by a
938 Standardized Single Disk Method. *American Journal of Clinical Pathology*, v. 45, p.493–
939 496, 1966.
- 940
941 BENGTSSON-PALME, J. et al. Using metagenomics to investigate human and
942 environmental resistomes, *Journal of Antimicrobial Chemotherapy*, v. 72, n. 10, p. 2690–
943 2703, 2017.
- 944
945 BONNET, R. et al. A novel CTX-M beta-lactamase (CTX-M-8) in cefotaxime-resistant
946 Enterobacteriaceae isolated in Brazil. *Antimicrobial Agents Chemotherapy*, v.44, n.7,
947 p.1936-42, Jul. 2000.
- 948
949 BOONYASIRI, A. et al. Prevalence of antibiotic resistant bacteria in healthy adults,
950 foods, food animals, and the environment in selected areas in Thailand. *Pathogens and*
951 *Global Health*, v. 108, p.235–45, 2014.
- 952
953 BUSH, K.; JACOBY, G. A. Update functional classification of beta-
954 lactamases. *Antimicrobial Agents Chemotherapy*, v. 54, n. 3, p. 969-76, 2010.
- 955
956 BRISOLA, M. C. et al. *Escherichia coli* used as a biomarker of antimicrobial resistance
957 in pig farms of Southern Brazil. *Science of the Total Environment*, v. 647, p. 362-368, 10
958 January 2019.

- 959
- 960 BROWER, C.H. et al. The prevalence of Extended-Spectrum Beta-Lactamase-Producing
961 Multidrug-Resistant *Escherichia coli* in Poultry Chickens and Variation According to
962 Farming Practices in Punjab, India. *Environmental Health Perspectives*, v.125, n.7,
963 077015, Jul. 20, 2017.
- 964
- 965 BRUNTON, L.A.; REEVES, H. E.; SNOW, L. C.; JONES, J. R. A. A longitudinal field
966 trial assessing the impact of feeding waste milk containing antibiotic residues on the
967 prevalence of ESBL-producing *Escherichia coli* in calves. *Preventive Veterinary
968 Medicine*, v.117, n.2, p.403-12, Nov. 15, 2014.
- 969
- 970 CANTÓN, R.; GONZÁLEZ-ALBA, J.M.; GALÁN, J.C. CTX-M Enzymes: Origin and
971 Diffusion. *Frontiers in Microbiology*, v.3, article 110, Apr. 2, 2012.
- 972
- 973 CARATTOLI, A. Animal reservoirs for extended spectrum beta-lactamase producers.
974 *Clinical Microbiology and Infection*. Jan; 14 Suppl 1:117-23, 2008.
- 975
- 976 CAREY, A. M. et al. Prevalence and Profiles of Antibiotic Resistance Genes *mph(A)* and
977 *qnrB* in Extended-Spectrum Beta-Lactamase (ESBL)- Producing *Escherichia coli*
978 Isolated from Dairy Calf Feces. *Microorganisms*, v. 10, p. 411. 2022.
- 979
- 980 CERGOLE-NOVELLA, M.C. et al. First description of *blaCTX-M-14* and *blaCTX-M-15*-
981 Producing *Escherichia coli* isolates in Brazil. *Microbial Drug Resistance*, v. 16, n. 3, p.
982 177-84, 2010.
- 983
- 984 COLOBATIU, L. et al. First description of plasmid-mediated quinolone resistance
985 determinants and beta-lactamase encoding genes in non-typhoidal *Salmonella* isolated
986 from humans, one companion animal and food in Romania. *Gut Pathogens*, v. 16, n. 07.
987 2015.
- 988
- 989 COLOM, K. et al. Simple and reliable multiplex PCR assay for detection of *blaTEM*,
990 *blaSHV* and *blaOXA-1* genes in Enterobacteriaceae. *FEMS Microbiology Letters*, v.223,
991 n.2, p-147-51, 2003.
- 992

- 993 COQUE, T. M.; BAQUERO, F.; CANTON, R. Increasing prevalence of ESBL-
994 producing Enterobacteriaceae in Europa. *Eurosurveillance*, v. 13, n.47, 19044, Nov. 20,
995 2008.
- 996
- 997 COURA, F. M. Longitudinal study of *Salmonella* spp., diarrheagenic *Escherichia coli*,
998 Rotavirus, and Coronavirus isolated from healthy and diarrheic calves in a Brazilian dairy
999 herd. *Tropical Animal Health Production*, v.47, p. 3-11, 2015.
- 1000
- 1001 CHONG, Y.; SHIMODA, S.; SHIMONO, N. Current epidemiology, genetic evolution,
1002 and clinical impact of extended-spectrum β -lactamase-producing *Escherichia coli* and
1003 *Klebsiella pneumoniae*. *Infection, Genetics and Evolution*, v.61, p.185–8, 2018.
- 1004
- 1005 CLSI, CLINICAL AND LABORATORY STANDARDS INSTITUTE. Performance
1006 Standards for Antimicrobial Susceptibility Testing. 31st ed. CLSI supplement M100, v.
1007 41, n. 03. USA, 2021.
- 1008
- 1009 EJAZ, H. et al. Molecular analysis of *blaSHV*, *blaTEM*, and *blaCTX-M* in extended-
1010 spectrum β -lactamase producing *Enterobacteriaceae* recovered from fecal specimens of
1011 animals. *PLoS ONE*, v. 16, n.01: e0245126, 2021.
- 1012
- 1013 ELLER, C. et al. Emergence of extended-spectrum β -lactamase (ESBL) CTX-M-8 in
1014 Germany. *Journal Antimicrobial Chemotherapy*, v.69, n.2, p.562-4, Feb. 2014.
- 1015
- 1016 ELSHAFIEE, E. A. et al. carbapenem-resistant *Pseudomonas aeruginosa* originating
1017 from farm animals and people in Egypt. *Journal Veterinary Research*, v. 63, 2019.
- 1018
- 1019 EWERS, C. et al. Emergence of human pandemic O25:H4-ST131 CTX-M-15 extended-
1020 spectrum-beta-lactamase-producing *Escherichia coli* among companion animals. *Journal*
1021 *Antimicrobial Chemotherapy*, v.65, n.4, p.651-60, Apr. 2010.
- 1022
- 1023 EWERS, C. et al. OXA-23 and ISAbal–OXA-66 class D β -lactamases in *Acinetobacter*
1024 *baumannii* isolates from companion animals. *International Journal of Antimicrobial*
1025 *Agents*, v. 49, n. 1, p.-37-44, 2017.
- 1026

- 1027 FERNANDES, M. R. et al., Zooanthroponotic Transmission of Drug-Resistant
1028 *Pseudomonas aeruginosa*, Brazil. *Emerging Infectious Diseases*. v, 24, n. 06, p. 1160-
1029 1162, 2018.
- 1030
- 1031 FRASAO, B. da SILVA et al. Detection of fluoroquinolone resistance by mutation in
1032 *gyrA* gene of *Campylobacter* spp. Isolates from broiler and laying (*Gallus gallus*
1033 *domesticus*) hens, from Rio de Janeiro State, Brazil. *Ciência Rural*, v. 45, n. 11, p. 2013-
1034 2018, nov, 2015.
- 1035
- 1036 GANGOUÉ-PIÉBOJI, J. et al. Extended-Spectrum-β-Lactamase-Producing
1037 *Enterobacteriaceae* in Yaounde, Cameroon. *Journal of Clinical Microbiology*, v. 43, n.
1038 07, 2005.
- 1039
- 1040 GUNDTRAN, R. S. et al. Prevalence and distribution of *blaCTX-M*, *blaSHV*, *blaTEM* genes in
1041 extended-spectrum β-Lactamase- producing *E. coli* isolates from broiler farms in the
1042 Philippines. *BMC Veterinary Research*, v.15, n.227, p.1-8, 2019.
- 1043
- 1044 HAN, R. et al. Elimination kinetics of ceftiofur hydrochloride in milk after an 8-day
1045 extended intramammary administration in healthy and infected cows. *PLoS One*, v.12,
1046 n.11, e0187261, nov. 2, 2017.
- 1047
- 1048 JARRIGE, N. et al. Effects of antimicrobial exposure on the antimicrobial resistance of
1049 *Escherichia coli* in the digestive flora of dairy calves. *Preventive Veterinary Medicine*, v.
1050 185:105177. Dec 2020.
- 1051
- 1052 JARLIER, V. et al. Extended broad-spectrum betalactamases conferring transferable
1053 resistance to newer beta-lactam agents in Enterobacteriaceae: hospital prevalence and
1054 susceptibility patterns. *Reviews Infection Disease*, v. 10, p. 867–878, 1988.
- 1055
- 1056 MAGIORAKOS, A. P.; SRINIVASAN, A.; CAREY, R. B. Multidrug-resistant,
1057 extensively drug-resistant, and pandrug-resistant bacteria: an international expert
1058 proposal for interim standard definitions for acquired resistance. *Clinical of Microbiology
1059 Infection*, v.18, n.3, p.268-281, 2011.
- 1060

- 1061 MELO, L. C. et al. Prevalence and molecular features of ESBL/pAmpC-producing
1062 Enterobacteriaceae in healthy and disease companion animals in Brazil. *Veterinary*
1063 *Microbiology*, v. 221, p. 59-66. 2018.
- 1064
- 1065 MUZAHEED, D. O. I. et al. High prevalence of CTX-M-15-producing *Klebsiella*
1066 *pneumoniae* among inpatients and outpatients with urinary tract infection in Southern
1067 India. *Journal Antimicrobial Chemotherapy*, v.61, n.6, p.1393-4, jun., 2008.
- 1068
- 1069 NESPOLO, N. M. et al. Ocorrência de *Escherichia coli* O157: H7 e O26 sorbitol
1070 negativas em matadouro frigorífico de bovino e suscetibilidade a antimicrobianos.
1071 *Arquivos do Instituto Biológico*, v. 81, n. 3, p. 209-217, 2014.
- 1072
- 1073 OIE, World Organization for Animal Health. Antimicrobial agents of veterinary
1074 importance at its 75th General Session in May 2007 (Resolution No. XXVIII). June 2021.
- 1075
- 1076 OLIVEIRA, C. J. B. et al. Methicillin-resistant *Staphylococcus aureus* from Brazilian
1077 dairy farms and identification of novel sequence types. *Zoonoses Public Health*, v. 63,
1078 p.97–105. 2016.
- 1079
- 1080 PALMEIRA, J.D., et al. Epidemic spread of Incl1/Pst113 plasmid carrying the Extended-
1081 Spectrum Beta-Lactamase (ESBL) *bla*CTX-M-8 gene in *Escherichia coli* of Brazilian
1082 catte. *Veterinary Microbiology*, v. 243, 108629, April 2020.
- 1083
- 1084 PATERSON, D. L.; BONOMO, R. A. Extended-spectrum beta-lactamases: a clinical
1085 update. *Clinical Microbiology Reviews*, v.8, n.4, p.657-86, 2005.
- 1086
- 1087 POIREL, L. et al. Antimicrobial resistance in *Escherichia coli*. *Microbiology Spectrum*
1088 *Journal*, v.6, n.4, ARBA-0026-2017. 2018.
- 1089
- 1090 POMBA, C. et al. First Report of OXA-23-Mediated Carbapenem Resistance in Sequence
1091 Type 2 Multidrug-Resistant *Acinetobacter baumannii* Associated with Urinary Tract
1092 Infection in a Cat. *Antimicrobial Agents and Chemotherapy*, vol. 58, n. 2, 2015.
- 1093

- 1094 PRIBUL, B. R. et al. Characteristics of quinolone resistance in *Salmonella* spp. Isolates
1095 from the food chain in Brazil. *Frontiers in Microbiology*, v. 8, p. 299. Mar 14, 2017.
- 1096
- 1097 RAHMAHANI, J. et al. Antimicrobial Resistance Profile of *Escherichia coli* From
1098 Cloacal Swab of Domestic Chicken in Surabaya Traditional Market. *Biochemical and*
1099 *Cellular Archives*, v, 20, n.1, p.2993-2997, 2020.
- 1100
- 1101 ROCHA, F.R.; PINTO, V.P.; BARBOSA, F.C. The Spread of CTX-M-Type Extended-
1102 Spectrum β -Lactamases in Brazil: A Systematic Review. *Microbial Drug Resistance*,
1103 v.22, n.4, p.301-11, Jun. 2016.
- 1104
- 1105 SCHMID, A.; HÖRMANSDORFER, S.; MESSELHÄUSSER, U. Prevalence of
1106 extended-spectrum β -lactamase-producing *Escherichia coli* on Bavarian dairy and beef
1107 cattle farms. *Applied and Environmental Microbiology*, v.79, n.9, p.3027-32, 2013.
- 1108
- 1109 SANTAJIT, S.; INDRAWATTANA, N. Mechanisms of Antimicrobial Resistance in
1110 ESKAPE Pathogens. *BioMed research international*, v. 2016, p.1-9, 2016.
- 1111
- 1112 SUKMAWINATA, E. et al. IncI1 Plasmid Associated with *bla*_{CTX-M-2} Transmission in
1113 ESBL-Producing *Escherichia coli* Isolated from Healthy Thoroughbred Racehorse,
1114 Japan. *Antibiotics*, v.9, n.70, p. 2-7, 2020.
- 1115
- 1116 SHAHRANI, M.; DEHKORDI, F. S.; MOMTAZ, H. Characterization of *Escherichia coli*
1117 virulence genes, pathotypes and antibiotic resistance properties in diarrheic calves in Iran.
1118 *Biological Research*, v. 47, n.1, p. 1-13, 2014.
- 1119
- 1120 SMITH T.; ORCUTT M. L. The bacteriology of the intestinal tract of young calves with
1121 special reference to the early diarrhea (“scours”). *Journal of Experimental Medicine*, v. 41,
1122 p.89–106, 1925.
- 1123
- 1124 TEKINER, L.; OZPINAR, H. Occurrence and characteristics of extended spectrum beta-
1125 lactamases-producing Enterobacteriaceae from foods of animal origin. *Brazilian Journal*
1126 *Microbiology*, v, 47, p.444-51, 2016.
- 1127

- 1128 TOLLENTINO, F.M. et al. High prevalence of bla(CTX-M) extended spectrum beta-
1129 lacramase genes in *Klebsiella pneumoniae* isolates from a tertiary care hospital: first
1130 report of bla(SHV-12), bla(SHV-31), bla(SHV-38), and bla(CTX-M-15) in Brazil.
1131 *Microbial Drug Resistance*, v.17, n.1, p.7-16, Mar. 2011.
- 1132
- 1133 TUTIJA, J. F et al. Molecular and phenotypic characterization of *Escherichia coli* from
1134 calves in an important meat-producing region in Brazil. *The Journal of Infection in*
1135 *Developing Countries*, v.16, n.6, p1030-1036. Jun 30, 2022.
- 1136
- 1137 WIDODO, A.; EFFENDI, M. H.; KHAIRULLAH, A.R. Extended-spectrum beta-
1138 lactamase (ESBL)-producing *Eschericia coli* from livestock. *Systematic Reviews in*
1139 *Pharmacy*, v. 11, n. 7, p. 382-392, 2020.
- 1140
- 1141 WHO, WORLD HEALTH ORGANIZATION. *Global Action Plan on Antimicrobial*
1142 *Resistance*. Geneva, Switzerland. World Health Organization, 2017.

1143 **5. (ARTIGO II)**

1144

1145 [Artigo formatado de acordo com as normas da revista *Journal of Biophotonics*]

1146

1147 **IDENTIFICATION OF MULTI-RESISTANT DIARRHEAGENIC *Escherichia***
1148 ***coli* BY USING FTIR SPECTROSCOPY AND MACHINE LEARNING: A**
1149 **FEASIBLE STRATEGY TO IMPROVE THE GROUP CLASSIFICATION**

1150

1151 Yasmin Garcia Marangoni¹, Thiago França², José Esteves², Ana Maranni², Cicero
1152 Cena^{2*}, Cassia R.B. Leal^{1*}

1153 *cicero.cena@ufms.br; *cassia.leal@ufms.br

1154

1155 1 UFMS – Universidade Federal de Mato Grosso do Sul, Faculdade de Medicina
1156 Veterinária e Zootecnia (FAMEZ), Campo Grande, MS, Brasil.

1157 2 UFMS –Universidade Federal de Mato Grosso do Sul, Optics and Photonic Lab
1158 (SISFOTON-UFMS), Campo Grande, MS, Brasil.

1159

1160 **ABSTRACT**

1161

1162 The identification of multidrug-resistant strains from *E. coli* species responsible for
1163 diarrhea in calves still faces many laboratory limitations and is necessary for properly
1164 monitoring the microorganism spread and control. Then, there is a need for the
1165 development of a screening tool for bacterial strain identification in microbiology
1166 laboratories, which must show easy implementation, fast response, and accurate results.
1167 The use of FTIR spectroscopy to identify microorganisms has been successfully
1168 demonstrated in the literature, including many bacterial strains, here we explored the
1169 FTIR potential for multidrug-resistant *E. coli* identification. First, we applied principal
1170 component analysis to observe the group formation tendency, the first results showed no
1171 clustering tendency with a messy sample score distribution, then we improved these
1172 results by properly selecting the main principal components. Finally, by using machine
1173 learning algorithms, a predicting model showed 70% overall accuracy in the validation
1174 test, which can be considered a great result for a screening tool. The present results

1175 demonstrate the viability of the method for microorganism identification, which can
1176 significantly contribute to its control.

1177 **Keywords:** multi-resistant bacteria, diarrheagenic, *Escherichia coli*, FTIR, machine
1178 learning.

1179

1180 INTRODUCTION

1181

1182 The introduction of antibiotics into clinical use was the greatest medical advance of
1183 the 20th century, and the rapid discovery of several classes of antibiotics in a relatively
1184 short period has led to the overuse of these drugs, which help promotes increasing
1185 antimicrobial resistance and antibiotics effectiveness loss [1]. The O'Neill report [2]
1186 predicted that without urgent action, ten million people a year will die from drug-resistant
1187 infections by 2050. As a warning to the world's population about the growing global
1188 resistance to antimicrobials, the WHO, World Health Organization [3] lists bacterial
1189 agents and their respective resistance to antimicrobials, divided into critical, high, and
1190 medium priority.

1191 Important strategies have been adopted in this scenario for world public health
1192 maintenance, between them the identification of multidrug-resistant bacteria in herds of
1193 animals [4], since we have observed the migration of this microorganism to a human-
1194 animal-environment interface, such as *Acinetobacter baumannii* [5, 6], *Pseudomonas*
1195 *aeruginosa* [7, 8], *Staphylococcus aureus* [9]; *Salmonellas species* [10, 11],
1196 *Campylobacter species* [12], and *Escherichia coli* [13, 14, 15].

1197 Besides, diarrhea in calves is one of the main causes of economic losses in Brazilian
1198 and worldwide livestock. Since the beginning of the 20th century, the worldwide
1199 scientific community points to the bacteria *E. coli* as one of the main agents involved in
1200 diarrhea of infectious origin; therefore, this bacterial genus and its pathogenic strains
1201 began to be studied and identified [16, 17, 18, 19].

1202 The *E. coli* bacteria is considered a commensal of the intestinal flora, only a small
1203 part of the strains has pathogenicity responsible for diseases and the differentiation of
1204 pathogenic (diarrheagenic) strains from non-pathogenic strains is based on the production
1205 of virulence factors [17]. Once *E. coli* bacteria with virulence factors spread in the cattle
1206 herd, serious concerns arise due to its capability of expressing enzymes that confer
1207 resistance to multiple antimicrobials [20, 17, 21]. Animal feces, which have multiresistant

1208 strains, serve as constant reservoirs for bacteria dissemination to humans [22, 23], the
1209 environment, and other animals [24, 25] which hinders the adequate therapy, increased
1210 morbidity, mortality, and cause economic losses to producers and industry [26, 27].

1211 Phenotypic and molecular methods have been used in bacterial culture to analyze the
1212 nucleic acid sequences and identify microorganisms with multidrug resistance
1213 characteristics, but their variable discriminatory capacity, high cost, and time-consuming
1214 experimental routine are difficult the implementation as a standard laboratory procedure
1215 [28, 29].

1216 In an attempt to overcome such limitations, several studies have been developed by
1217 using Fourier Transform Infrared Spectroscopy (FTIR) as a screening test [30, 31], which
1218 can obtain information about the sample chemical compounds thought their vibrational
1219 molecular modes, enabling the identification of sample molecular composition (nucleic
1220 acids, proteins, lipids, and carbohydrates) of these bacterial cells [32, 33, 34].

1221 The potential use of FTIR spectroscopy to discriminate, classify and identify
1222 microorganisms has been successfully demonstrated in the literature, including many
1223 bacterial strains from Gram-positive, and Gram-negative species [32, 35, 36, 37, 38, 39,
1224 40, 41]. The specific spectral patterns observed for each molecular group can be revealed
1225 with multivariate analysis and/or machine learning algorithms aid, which enable FTIR
1226 spectroscopy as an alternative for rapid bacterial identification at the subspecies level [42,
1227 43]. Despite its great applicability for microorganism identification in recent years, this
1228 technique's potential in microbiology routines is still underestimated, due to the difficult
1229 acceptance and comprehension of the methodology by the microbiological community
1230 [31].

1231 Therefore, here we explore the use of FTIR spectroscopy associated with machine
1232 learning algorithms for data analysis to identify and classify multidrug-resistant strains
1233 from *E. coli* species responsible for diarrhea in calves, which may add value to the
1234 microbiology laboratory toolbox and society [26, 27, 31]. Our study is focused on the
1235 development of a simple laboratory routine to sample preparation and data acquisition,
1236 then our data analysis aims to improve the method's accuracy for future implementation
1237 as a screening tool in bacterial strain identification in microbiology laboratories.

1238

1239

METHODS


1240

1241 **Sample preparation and FTIR spectra acquisition**

1242 A total of 80 *Escherichia coli* isolates were analyzed. The sample pool was
 1243 divided into 40 multidrug-resistant (MR) *E. coli* isolates and 40 replicates of *E. coli*
 1244 ATCC® 25922. First, they were screened for analysis and placed in Brain Heart Infusion
 1245 (BHI) broth for activation, subsequently, seeded on MacConkey Agar to verify purity and
 1246 isolate selected colonies. Then, the colonies of MR *E. coli* and ATCC *E. coli* were
 1247 suspended in 1 mL of BHI broth, and a resulting turbid inoculum was adjusted to the 0.5
 1248 McFarland scale (10^8 CFU/mL) before the measurements.

1249 A small aliquot (30 μ L) of the bacterial isolate was carefully deposited onto a flat
 1250 silicon substrate (SiO_2) by casting, followed by drying at 50°C per 30 minutes – this
 1251 procedure was repeated three times until the obtention of a thick film. Each sample was
 1252 produced in duplicate, and the samples were analyzed in two different spots from the
 1253 center to the border, to avoid inhomogeneity in the sample composition due to the drying
 1254 process [44]. Finally, the mid-infrared spectra were obtained by using an attenuated total
 1255 reflectance accessory (ATR) at Fourier Transform Infrared Spectrophotometer (Spectrum
 1256 100, Perkin Elmer). The spectra were collected from 4000 to 700 cm^{-1} , with 4 cm^{-1}
 1257 resolution and 10 scans. This entire experimental roadmap is illustrated in Figure 1.

1258



1259

1260 Figure 1: Experimental scheme describing the main steps from the sample (*E. coli*)
 1261 preparation to FTIR spectra acquisition.
 1262

1263 ***Data analysis and sample classification***

1264 The data analysis was performed in Python (version 3.9.12), using the Scikit-learn
 1265 package (version 1.1.2) [45]. First, the FTIR spectra from two different spots of duplicate
 1266 samples were averaged and then subjected to Standard Normal Variate (SNV) pre-
 1267 processing method, which removes the variation from the baseline and rescales the
 1268 spectral intensity, to prevent interference in the data analysis due to random experimental
 1269 variations [46].

1270 The average FTIR-SNV spectra for MR *E. coli* and ATCC *E. coli* group, in the
 1271 4000 to 800 cm⁻¹ range, were submitted to principal component analysis (PCA) [47]. PCA
 1272 is an unsupervised method that will project our pre-processed data set into a new
 1273 dimension (PCs – principal components) which aims to maximize the data variance, this
 1274 dimensionally reduced data set retains most of the information from the original variables
 1275 and shows how each data sample is distributed in this new dimension (score plot),
 1276 allowing cluster the similar samples and distinguish groups. Each PC represents a
 1277 percentage of the data variance and, allows us to analyze the main spectral range that most
 1278 contributes to the data variance percentage through the loading plot. PCA is an important
 1279 step to visualize the group classification tendency. Here we analyzed three different
 1280 ranges: (i) 4000 to 800 cm⁻¹; (ii) 3000 to 2800 cm⁻¹, and (iii) 1800 to 800 cm⁻¹, to use only
 1281 those vibrational modes that improve the group clustering and classification and eliminate
 1282 highly correlated data [48].

1283 The sample classification is performed by prediction models built by machine
 1284 learning (ML) algorithms using PC output data from 70% sample set. Before sample
 1285 classification tests, we must determine the ideal number of PCs used by ML algorithms
 1286 to avoid overfitting and underfitting [49, 50]. Here we used the Feature Selection
 1287 Recursive Feature Elimination (RFE), which selects the main PCs that most contribute to
 1288 achieving high accuracy and remove other PCs with the weakest contribution to correct
 1289 sample classification in ML tests [51]. The use or removal of a determined PC was made
 1290 based on the accuracy achieved by using Linear Discriminant Analysis (LDA) to classify
 1291 the samples in a Leave One Out Cross-validation (LOOCV) test.

1292 In a brief description, Discriminant Analysis (DA) classifies the sample based on
 1293 the distance between the sample data and the contour built by using a linear (L) or

1294 quadratic (Q) function to separate the classes (group) [52]. In LOOCV, one sample is
1295 taken from the data set, and the others are used to build the prediction model (training).
1296 Then, the prediction model accuracy is tested by using the sample data withdrawal from
1297 the data set. The procedure is repeated until all sample data have been tested [53].

1298 After determining the ideal number of PCs, and which PCs most contribute to
1299 sample classification in each spectral range analyzed, a LOOCV test was performed – by
1300 using the respective RFE-PCs data for each range – based on three different methods: (i)
1301 DA (described above); (ii) k-Nearest Neighbor (KNN), which uses the Euclidean distance
1302 between k closest neighbors to classify the sample [54]; and (iii) Support Vector Machine
1303 (SVM), which organizes each sample class through the optimization of a hyperplane –
1304 the hyperplane can be linear or nonlinear, being optimized to reach high performance –
1305 between the classes [55]. Finally, we were able to determine the best spectral range, ML
1306 algorithm, and PCs to build a predicting model, whose ability for generalization was
1307 tested in an external validation test by using 30% of the sample set.


1308

1309 **RESULTS AND DISCUSSION**

1310

1311 The average FTIR-SNV spectra for *E. coli* ATCC, and MR *E. coli* samples, Figure
1312 2, exhibit great similarity between them, with a small standard deviation into the data set.
1313 It suggests that direct identification of the isolate is not possible, and the data set shows
1314 great coherence. There is a remarkable presence of the three most pronounced bands in
1315 the spectra, the first around 1600 cm⁻¹ is the result of overlapped bands 1639 and 1583
1316 cm⁻¹, assigned to Amides I and II from proteins, which is relatively wide, suggesting the
1317 presence of a third band, mainly due to the small shoulder observed around 1500 cm⁻¹.
1318 The vibrational bands assigned to Amides I (N-H and C=O) and II (N-H and C-N) have
1319 shown great contributions to biological sample classification in the literature [56, 57].
1320 The second band around 1400 cm⁻¹ can be assigned to C-H and C=O vibrational modes
1321 from lipids and proteins, while the third band around 1080 cm⁻¹ is usually assigned to
1322 nucleic acid and phospholipids. Also, weak bands assigned to C-H, and C-H-O vibrational
1323 modes from carbohydrate below 1000 cm⁻¹ was identified in the spectra [58, 59].

1324



1326 **Figure 2:** Average FTIR-SNV spectra for (a) *E. coli* ATCC (blue line), and (b) multidrug-resistant
 1327 (MR) *E. coli* isolates. The main vibrational assignments are indicated in detail and their
 1328 related molecular groups (proteins, lipids, carbohydrates, and fatty acids) are identified by colors.
 1329 The symbol δ indicates the deformation vibrations.

1330 Figure 3 shows the principal component analysis results for the FTIR-SNV
 1331 spectral data from *E. coli* ATCC, and MR *E. coli* sample groups. On the left side, we can
 1332 observe the score plot, and on the right side, the loading plot for (i) 4000 to 800 cm⁻¹; (ii)
 1333 3000 to 2800 cm⁻¹, and (iii) 1800 to 800 cm⁻¹ range. For all cases analyzed two main
 1334 characteristics stand out, the score plot exhibits a single cluster for both groups, which
 1335 hinders the future group classification, and the loading shows few regions with a small
 1336 contribution to the data variance, suggesting a small contribution of the bands for data
 1337 variance in PC1 and PC2. We found PC1 and PC2 responsible for 79.2%, 96.8%, and
 1338 74.9% of data variance at 4000 to 800 cm⁻¹, 3000 to 2800 cm⁻¹, and 1800 to 800 cm⁻¹
 1339 range, respectively, besides this great contribution for data variance previous studies have
 1340 shown that the first PCs can be ignored, and high order PCs can be used to improve the
 1341 group classification for ML algorithms [60].

1343




Figure 3: Principal Component Analysis for *E. coli* ATCC (blue circles/stars), and multidrug-resistant (MR) *E. coli* (red circles/stars) isolates average FTIR-SNV spectra. The score plot and respective loading for the three different ranges were analyzed: (a-b) 4000 to 800 cm^{-1} ; (c-d) 3000 to 2800 cm^{-1} ; and (e-f) 1800 to 800 cm^{-1} . The circles represent 70% of the sample data set used to build the predicting model, and the stars represent 30% of the sample data set used for external validation.

Usually, the proper choice of spectral range helps to improve group classification and clustering formation [61], since we use only spectral information that most contributes to clustering from those with highly correlated data, which hinders cluster formation. But here we couldn't succeed with this strategy probably because of the high similarities between the groups involved, and the highly correlated data present. The presence of antibiotic resistance genes in the MR *E. coli* group was mainly identified as resulting of the ESBL enzyme – previous study [62] - which is responsible for amine

1361 beta-lactam ring hydrolysis, resulting in antibiotics inactivation on the bacterial cell walls.
1362 Since it represents a break in the C-N bond (Amide group) and the appearance of N-H
1363 and O-H bonding, almost no difference in the IR spectra will be observed. Because, the
1364 C-N and N-H vibrational modes are superimposed around 1580 cm^{-1} , and O-H vibrational
1365 mode, around 3278 cm^{-1} , is too wide to provide enough information for group
1366 differentiation.

1367 Then, an alternative to improve group clustering and separation is the use of
1368 Feature Selection Recursive Feature Elimination (RFE) to find the best PCs (data
1369 projection) that most contribute to data separation. The RFE can select PCs that most
1370 contribute to achieving high accuracy and remove those with a small contribution. Here,
1371 the main PCs were determined by the overall accuracy achieved in a Leave One Out
1372 Cross-validation (LOOCV) test by using Linear Discriminant Analysis (LDA) [60].The
1373 main PCs found for our data were: PC2, PC3, PC7, PC8, and PC19 for the 4000 to 800
1374 cm^{-1} range, responsible for 20.26% of data variance; PC5, PC13, PC21, PC36, and PC46
1375 for the 3000 to 2800 cm^{-1} range, responsible for 0.49% of data variance; and PC4, PC6,
1376 PC7, PC12, PC13, PC17, PC28, PC33, and PC48 for the 1800 to 800 cm^{-1} range
1377 responsible for 8.58% of data variance.

1378 Figure 4 shows the violin plot for the 9 most relevant PCs, in 1800 to 800 cm^{-1}
1379 range, with a clear monomodal distribution for the score data project over its respective
1380 PC for both sample groups, except for PC7, which exhibits a wide distribution for MR *E.*
1381 *coli* group, and a double peak maximum for *E. coli* ATCC, similar behavior observed for
1382 MR *E. coli* group in PC6. But the more important characteristic to be observed is the
1383 median value (dashed thick line around the peak center), which assumes a better distinct
1384 position projection over the axis for these PCs compared to the others and improves the
1385 clustering and group classification of our data in the LOOCV test with LDA.

1386




Figure 4: Violin plot for nine main PCs normalized scores selected by RFE in 1800 to 800 cm^{-1} range from *E. coli* ATCC® 25922 (blue), and multidrug-resistant (MR) *E. coli* isolates. The thick dashed line represents the data distribution median. The data variance percentage of each PC is described above each plot.

Figure 5 exemplifies the importance of these RFE-PCs. The score plot and loading for PC4, PC17, and PC28 demonstrate the improvement of data clustering and group separation. They were responsible for only 5.05% of the data variance, similar results have been reported in the literature [60]. Besides that, the loading plot for each PC, Figure 5 (right column), shows a contribution for data variance in the 1600 to 1000 cm^{-1} range in accordance with the main bands identified in the FTIR-SNV spectra (Figure 2) assigned to protein and phospholipids molecules. In this case, with great similarity among the samples, then between the spectra – high data correlation – the first PCs can be responsible for a great percentage of data variance, with a small contribution to group separation. However, low data variance PCs can still be useful for group classification and better influence the algorithm performance, as demonstrated by Figure 5 results.




Figure 5: (a) score plot clustering improvement due to PCs selection by RFE., and (b) loading for PC4, PC17, and PC28 from *E. coli* ATCC® 25922 (blue square), and (b) multidrug-resistant (MR) *E. coli* (red circles) isolates.

The RFE-PCs were submitted to machine learning algorithms (DA, SVM, and KNN) to build a prediction model for sample classification, the overall accuracy achieved for each model by using different functions for classification in the LOOCV test is summarized in Figure 6 (a). The maximum overall accuracy achieved by the predicting models was 85% for Linear SVM by using 9 PCs – responsible for 8.59% of data variance – within the 1800 to 800 cm⁻¹ range, and 5 PCs in the 4000 to 800 cm⁻¹ range. The Weighted KNN also achieved an overall accuracy of 85% in the 4000 to 800 cm⁻¹ range with 5 PCs, which was responsible for 20.26% of data variance.




Figure 6: (a) Overall accuracy obtained in the LOOCV tests for Discriminant analysis (DA), K-nearest neighbor (KNN), and Support vector machine (SVM) algorithms. Different ranges were

1421 analyzed with respective RFE-PCs: (i) 4000 to 800 cm⁻¹ (gray bar color with right inclined line
 1422 pattern) using 5 PCs; (ii) 3000 to 2800 cm⁻¹ (light blue color with horizontal line pattern) by using
 1423 5 PCs; (iii) 1800 to 800 cm⁻¹ (purple color with left inclined line pattern) by using 9 PCs. The
 1424 highest overall accuracy was 85% for Linear SVM at (4000 to 800 cm⁻¹) and (1800 to 800 cm⁻¹)
 1425 ranges, and Weighted KNN at (4000 to 800 cm⁻¹) range. Confusion matrix for the performance of
 1426 Linear SVM algorithm in 1800 to 800 cm⁻¹ range by using 9 PCs (b) LOOCV test with 70% data
 1427 set, and (c) external validation test with 30% data set. The input PCA data from average FTIR-
 1428 SNV spectra of *E. coli* ATCC® 25922, and multidrug-resistant (MR) *E. coli* isolates.
 1429

1430 Besides both models achieving the same overall accuracy of 85% in the 4000 to
 1431 800 cm⁻¹ range, the W-KNN and L-SVM showed a much lower generalizability of the
 1432 model in the external validation test, when compared to L-SVM in the 1800 to 800 cm⁻¹
 1433 range. The external validation overall accuracy for L-SVM was 55%, and only 50% for
 1434 W-KNN (see Supp. Mat.). However, the L-SVM with 85% of overall accuracy in the
 1435 1800 to 800 cm⁻¹ range, achieved 90% and 80% of sensitivity and specificity,
 1436 respectively, Figure 6(b). Then, in the external validation test, Figure 6(c), the overall
 1437 accuracy decreased to 70%, and achieved 90% and 50% of sensitivity and specificity,
 1438 respectively. As can be observed in Figure 6(c), the prediction model failed to identify *E.*
 1439 *coli* ATCC samples, which is less problematic for a trial method considering the need for
 1440 large-scale tests to a fast decision in livestock management.

1441 The current study presented an easy and suitable methodology for the
 1442 identification of MR-*E. coli* bacteria by FTIR spectroscopy and machine learning
 1443 algorithms but must keep in mind that the methodology and results can be still improved,
 1444 so the possibility for new studies is open. Here, we discussed the FTIR limitations to
 1445 evidenced spectral changes due role played by ESBL enzyme to create MR bacteria, so
 1446 new photonic techniques must be explored for better data acquisition. In our study, we
 1447 explore a promising route to build a prediction model with good performance in the
 1448 external validation test, but we can still search for the best hyperparameters for the
 1449 algorithm or even explore new pre-processing data methods and algorithms to build the
 1450 prediction model and improve its generalization capacity in the external validation test.
 1451

1452 CONCLUSIONS

1453 The FTIR spectra, in the 1800 to 800 cm⁻¹ range, of multi-resistant diarrheagenic
 1454 *Escherichia coli*, isolates obtained from calves' feces showed great potential for
 1455 microorganism identification. The usual principal component analysis was not able to
 1456 provide a promising clustering formation for future sample classification. Then, we

1458 applied the Feature Selection Recursive Feature Elimination (RFE) algorithm, from
 1459 which the most important PCs were chosen before being used in machine learning
 1460 algorithms. The score plot of PC4 x PC17 x PC28 clearly demonstrated the improvement
 1461 of clustering tendency and group separation, here 9 PCs were selected by RFE and used
 1462 to build a prediction model by using Linear SVM. The prediction model showed an 85%
 1463 overall accuracy in the LOOCV test, and achieved 70% overall accuracy, with 90%
 1464 sensitivity and 50% specificity in the external validation test.

1465

1466 ACKNOWLEDGMENTS

1467

1468 The present study was carried out with the support of the Universidade Federal de
 1469 Mato Grosso do Sul – UFMS/MEC, Brazil, Coordenação de Aperfeiçoamento de Pessoal
 1470 de Nível Superior (CAPES), code 001. Conselho Nacional de Desenvolvimento
 1471 Científico e Tecnológico (CNPQ), code 403651/2020-5; 302525/2022-0; 440214/2021-
 1472 1. Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia do Estado
 1473 de Mato Grosso do Sul (FUNDECT), code 007/2019; 360/2022.

1474

1475 REFERENCES

1476

- 1477 [1] KATZ, L.; BALTZ, R. H. Natural product discovery: past, present, and future. *Journal of Industrial Microbiology and Biotechnology*, v. 43, p. 155-176, 2016.
- 1478 [2] O'NEILL, J. *Tackling drug-resistant infections globally: final report and recommendations. Review on Antimicrobial Resistance*. Wellcome Trust and HM Government. Disponível em:
https://amrreview.org/sites/default/files/160525_Final%20paper_with%20cover.pdf.
 1483 2016.
- 1484 [3] WHO, WORLD HEALTH ORGANIZATION. *Global Action Plan on Antimicrobial Resistance*. Geneva, Switzerland. World Health Organization, 2017.
- 1485 [4] HAN, R. et al. Elimination kinetics of ceftiofur hydrochloride in milk after an 8-day extended intramammary administration in healthy and infected cows. *PLoS One*, v.12, n.11, e0187261, nov. 2, 2017.
- 1486 [5] POMBA, C. et al. First Report of OXA-23-Mediated Carbapenem Resistance in Sequence Type 2 Multidrug-Resistant *Acinetobacter baumannii* Associated with Urinary Tract Infection in a Cat. *Antimicrobial Agents and Chemotherapy*, vol. 58, n. 2, 2015.
- 1487 [6] EWERS, C. et al. OXA-23 and ISAbal-OXA-66 class D β -lactamases in *Acinetobacter baumannii* isolates from companion animals. *International Journal of Antimicrobial Agents*, v. 49, n. 1, p.-37-44, 2017.
- 1488 [7] FERNANDES, M. R. et al., Zooanthroponotic Transmission of Drug-Resistant *Pseudomonas aeruginosa*, Brazil. *Emerging Infections Diseases*. v, 24, n. 06, p. 1160-1162, 2018.

- 1498 [8] ELSHAFIEE, E. A. et al. carbapenem-resistant *Pseudomonas aeruginosa* originating
1499 from farm animals and people in Egypt. *Jornal Veterinary Reseasch*, v. 63, 2019.
1500 [9] OLIVEIRA, C. J. B. et al. Methicillin-resistant *Staphylococcus aureus* from Brazilian
1501 dairy farms and identification of novel sequence types. *Zoonoses Public Health*, v. 63,
1502 p.97–105. 2016.
1503 [10] COLOBATIU, L. et al. First description of plasmid-mediated quinolone resistance
1504 determinants and beta-lactamase encoding genes in non-typhoidal *Salmonella* isolated
1505 from humans, one companion animal and food in Romania. *Gut Pathogens*, v. 16, n. 07.
1506 2015.
1507 [11] PRIBUL, B. R. et al. Characteristics of quinolone resistance in *Salmonella* spp.
1508 Isolates from the food chain in Brazil. *Frontiers in Microbiology*, v. 8, p. 299. Mar 14,
1509 2017.
1510 [12] FRASAO, B. da SILVA et al. Detection of fluoroquinolone resistance by mutation
1511 in *gyrA* gene of *Campylobacter* spp. Isolates from broiler and laying (*Gallus gallus*
1512 *domesticus*) hens, from Rio de Janeiro State, Brazil. *Ciência Rural*, v. 45, n. 11, p. 2013-
1513 2018, nov, 2015.
1514 [13] MELO, L. C. et al. Prevalence and molecular features of ESBL/pAmpC-producing
1515 Enterobacteriaceae in healthy and disease companion animals in Brazil. *Veterinary*
1516 *Microbiology*, v. 221, p. 59-66. 2018.
1517 [14] BRISOLA, M. C. et al. *Escherichia coli* used as a biomarker of antimicrobial
1518 resistance in pig farms of Southern Brazil. *Science of the Total Environment*, v. 647, p.
1519 362-368, 10 January 2019.
1520 [15] PALMEIRA, J.D., et al. Epidemic spread of Incl1/Pst113 plasmid carrying the
1521 Extended-Spectrum Beta-Lactamase (ESBL) *blaCTX-M-8* gene in *Escherichia coli* of
1522 Brazilian catte. *Veterinary Microbiology*, v. 243, 108629, April 2020.
1523 [16] SMITH T.; ORCUTT M. L. The bacteriology of the intestinal tract of young calves
1524 with special referenceto the early diarrhea (“scours”). *Journal of Experimental Medicine*,
1525 v. 41, p.89–106, 1925.
1526 [17] COURIA, F. M. Longitudinal study of *Salmonella* spp., diarrheagenic *Escherichia coli*,
1527 Rotavirus, and Coronavirus isolated from healthy and diarrheic calves in a Brazilian
1528 dairy herd. *Tropical Animal Health Production*, v.47, p. 3-11, 2015.
1529 [18] CRUVINEL, L. B. et al. Surtos consecutivos ocasionados por *Eimeria zuernii* em
1530 bezerros de corte de uma propriedade do estado de São Paulo. *Pesquisa Veterinária*
1531 *Brasileira*, v.38, n. 2, p.277-284, fevereiro, 2018.
1532 [19] TUTIJA, J. F et al. Molecular and phenotypic characterization of *Escherichia coli*
1533 from calves in an important meat-producing region in Brazil. *The Journal of Infection in*
1534 *Developing Countries*, v.16, n.6, p1030-1036. Jun 30, 2022.
1535 [20] BLANCHARD, P.C. 2012. Diagnostics of dairy and beef cattle diarrhea. *Veterinary*
1536 *Clinics of North America: Food Animal Practice*, v.28, p443-464, 2012.
1537 [21] POIREL, L. et al. Antimicrobial resistance in *Escherichia coli*. *Microbiology*
1538 *Spectrum Journal*, v.6, n.4, ARBA-0026-2017. 2018.
1539 [22] CARATTOLI, A. Animal reservoirs for extended spectrum beta-lactamase
1540 producers. *Clinical Microbiology and Infection*, Jan; 14 Suppl 1:117-23, 2008.
1541 [23] SCHMID, A.; HÖRMANSDORFER, S.; MESSELHÄUSSER, U. Prevalence of
1542 extended-spectrum β-lactamase-producing *Escherichia coli* on Bavarian dairy and beef
1543 cattle farms. *Applied and Environmental Microbiology*, v.79, n.9, p.3027-32, 2013.
1544 [24] BOONYASIRI, A. et al. Prevalence of antibiotic resistant bacteria in healthy adults,
1545 foods, food animals, and the environment in selected areas in Thailand. *Pathogens and*
1546 *Global Health*, v. 108, p.235–45, 2014.

- 1547 [25] TEKINER, L.; OZPINAR, H. Occurrence and characteristics of extended spectrum
1548 beta-lactamases-producing Enterobacteriaceae from foods of animal origin. *Brazilian*
1549 *Journal of Microbiology*, v. 47, p.444-51, 2016.
- 1550 [26] SANTAJIT, S.; INDRAWATTANA, N. Mechanisms of Antimicrobial Resistance
1551 in ESKAPE Pathogens. *BioMed research international*, v. 2016, p.1-9, 2016.
- 1552 [27] CHONG, Y.; SHIMODA, S.; SHIMONO, N. Current epidemiology, genetic
1553 evolution, and clinical impact of extended-spectrum β -lactamase-producing Escherichia
1554 coli and Klebsiella pneumoniae. *Infection, Genetic and Evolution*, v.61, p.185–8, 2018.
- 1555 [28] JARLIER, V. et al. Extended broad-spectrum betalactamases conferring transferable
1556 resistance to newer beta-lactam agents in Enterobacteriaceae: hospital prevalence and
1557 susceptibility patterns. *Reviews of Infectious Diseases*, v. 10, p. 867–878, 1988.
- 1558 [29] SABAT, A. J.; BUDIMIR, A.; NASHEV, D. et al. Overview of molecular typing
1559 methods for outbreak detection and epidemiological surveillance. *Eurosurveillance*, v.
1560 18, n. 4, 2013.
- 1561 [30] MAQUELIN, K. et al. Identification of medically relevant microorganisms by
1562 vibrational spectroscopy. *Journal of Microbiological Methods*, v.51, p. 255–271, 2002.
- 1563 [31] NOVAIS, A.; FREITAS, A. R.; RODRIGUES, C. et al. 27Fourier transform infrared
1564 spectroscopy: unlocking fundamentals and prospects for bacterial strain typing. *European*
1565 *Journal of Clinical Microbiology & Infection Diseases*, v.38, n.3, p.427-448, 2019.
- 1566 [32] HELM, D.; LABISCHINSKI, H.; SCHALLEHN, G. et al. Classification and
1567 identification of bacteria by Fourier-transform infrared spectroscopy. *Journal of General*
1568 *Microbiology*, v. 137, p. 69-79, 1991.
- 1569 [33] SETTLE, F. A. *Handbook of Instrumental Techniques for Analytical Chemistry*.
1570 Prentice Hall. New Jersey. USA.1997.
- 1571 [34] STUART, B. *Infrared Spectroscopy: Fundamentals and Applications*. John Wiley
1572 & Sons. West Sussex. England. 2004.
- 1573 [35] MARIEY, L. et al. Discrimination, classification, identification of microorganisms
1574 using FTIR spectroscopy and chemometrics. *Vibrational Spectroscopy*, v.26, n.2, p.151-
1575 159, 2001.
- 1576 [36] BEUTIN, L.; WANG, Q.; NAUMANN, D. et al. Relationship between O-antigen
1577 subtypes, bacterial surface structures and O-antigen gene clusters in *Escherichia coli*
1578 O123 strains carrying genes for Shiga toxins and intimin. *Journal Medical of*
1579 *Microbiology*, v. 56, p.177–184, 2007.
- 1580 [37] COLABELLA C. et al. Merging FTIR and NGS for simultaneous phenotypic and
1581 genotypic identification of pathogenic Candida species. *PLoS ONE*, v. 12, n.12:
1582 e0188104, 2017.
- 1583 [38] POTOCKI, L. et al. FTIR and Raman Spectroscopy-Based Biochemical Profiling
1584 Reflects Genomic Diversity of Clinical Candida Isolates That May Be Useful for
1585 Diagnosis and Targeted Therapy of Candidiasis. *International Journal of Molecular*
1586 *Sciences*, v. 20, 988, 2019.
- 1587 [39] RUSTAM, R. G. et al. Discrimination of *Staphylococcus aureus* Strains from
1588 Coagulase-Negative Staphylococci and Other Pathogens by Fourier Transform Infrared
1589 Spectroscopy. *Analytical Chemistry*, v. 92, n. 7, p.4943-4948, 2020.
- 1590 [40] PEBOTUWA, S. et al. Influence of the Sample Preparation Method in
1591 Discriminating Candida spp. Using ATR-FTIR Spectroscopy. *Molecules*, v, 25, 1551,
1592 2020.
- 1593 [41] CORDOVANA, M. et al. Classification of *Salmonella enterica* of the (Para-)Typhoid
1594 Fever Group by Fourier-Transform Infrared (FTIR) Spectroscopy. *Microorganisms*, v. 9,
1595 853, 2021.

- 1596 [42] MEYERS, R. A. Encyclopedia of Analytical Chemistry (Applications, Theory, and
1597 Instrumentation). *Infrared Spectroscopy in Microbiology.*, p. 1–32, 2006.
1598 [43] SHI, H. et al. The strategy for correcting interference from water in Fourier transform
1599 infrared spectrum based bacterial typing. *Talanta*, v. 208, 2020.
1600 [44] CAMERON, J. M. et al. Biofluid spectroscopic disease diagnostics: a review on the
1601 processes and spectral impact of drying. *Journal of Biophotonics*, v. 11: e201700299,
1602 2018.
1603 [45] PEDREGOSA, F. et al. Scikit-learn: Machine learning in Python. *The Journal of
1604 Machine Learning Research*, v. 12, p. 2825-2830, 2011.
1605 [46] RINNAN, Å.; VAN DEN BERG, F.; ENGELSEN, S. B. Review of the most
1606 common pre-processing techniques for near-infrared spectra. *TrAC Trends in Analytical
1607 Chemistry*, v. 28, n. 10, p. 1201-1222, 2009.
1608 [47] JOLLIFFE, I. T.; CADIMA, J. Principal component analysis: a review and recent
1609 developments. *Philosophical Transactions of the Royal Society A: Mathematical,
1610 Physical and Engineering Sciences*, v. 374, n. 2065, p. 20150202, 2016.
1611 [48] CASARIL, A. E. Intraspecific differentiation of sandflies specimens by optical
1612 spectroscopy and multivariate analysis, *Journal of Biophotonic.*, v.14: e202000412 v.,
1613 2021.
1614 [49] GOMES, T. R. FTIR spectroscopy with machine learning: A new approach to animal
1615 DNA polymorphism screening. *Spectrochimica Acta Part A: Molecular and
1616 Biomolecular Spectroscopy*, 2021.
1617 [50] HASBI, N. H. et al. Pattern recognition for human diseases classification in spectral
1618 analysis, *Computation*, v. 10, p. 96, 2022.
1619 [51] THEERTHAGIRI, P. Predictive analysis of cardiovascular disease using gradient
1620 boosting-based learning and recursive feature elimination technique. *Intelligent Systems
1621 with Applications*, v. 16, p. 200121, 2022.
1622 [52] WU, W. et al. Comparison of regularized discriminant analysis linear discriminant
1623 analysis and quadratic discriminant analysis applied to NIR data. *Analytica Chimica Acta*,
1624 v. 329, n. 3, p. 257-265, 1996.
1625 [53] WONG, T. Performance evaluation of classification algorithms by k-fold and leave-
1626 one-out cross-validation. *Pattern Recognition*, v. 48, n. 9, p. 2839-2846, 2015.
1627 [54] MUCHERINO, A.; PAPAJORGJI, P. J.; PARDALOS, P. M. K-nearest neighbor
1628 classification. In: *Data mining in agriculture*. Springer, New York, NY, 2009. p. 83-106.
1629 [55] NOBLE, William S. What is a support vector machine? *Nature biotechnology*, v. 24,
1630 n. 12, p. 1565-1567, 2006.
1631 [56] LARIOS, G. et al. A new strategy for canine visceral leishmaniasis diagnosis based
1632 on spectroscopy and machine learning, *Journal of Biophotonics.*, v.11: e202100141 vp.,
1633 2021.
1634 [57] BRITO, E.C.A. Paracoccidioidomycosis screening diagnosis by FTIR spectroscopy
1635 and multivariate analysis, *Photodiagnosis and Photodynamic Therapy*, v.39, 102921,
1636 2022.
1637 [58] LEGAL, J. M. Applications of FTIR spectroscopy in structural studies of cells and
1638 bacteria. *Journal of Molecular Structure*, v. 242, p.397-407 Elsevier Science Publishers
1639 B.V., Amsterdam. 1991.
1640 [59] MARIETY, L. Discrimination, classification, identification of microorganisms using
1641 FTIR spectroscopy and chemometrics. *Vibrational Spectroscopy*, v. 26, p.151-159, 2021.
1642 [60] COELHO, M. L. et al. Canine visceral leishmaniasis diagnosis by UV spectroscopy
1643 of blood serum and machine learning algorithms. *Photodiagnosis and Photodynamic
1644 Therapy* (corrected proofs).

- 1645 [61] FRANCA, T; GONÇALVES, D; CENA, C. ATR-FTIR spectroscopy combined with
1646 machine learning for classification of PVA/PVP blends in low concentration. *Vibrational*
1647 *Spectroscopy*, v. 120: 103378, May 2022.
- 1648 [62] MARANGONI-GHOREYSHI, Y.G. et al. Identificação fenotípica e genotípica de
1649 *Escherichia coli* produtoras de beta-lactamases de espectro estendido (ESBL) isoladas de
1650 fezes de bezerros (pre-proofs).